

Crosstalk between the androgen and estrogen receptors in breast cancer

Easter Ndlovu

*Dissertation presented in fulfilment of the requirements for the degree
Masters of Science in Biochemistry
At the University of Stellenbosch*



Promoter: Dr. Donita Africander

Co-promoter: Prof. Ann Louw

March 2016

DECLARATION

By submitting this dissertation electronically, I declare that the entirety of the work contained therein is my own, original work, that I am the owner of the copyright thereof (unless to the extent explicitly otherwise stated) and that I have not previously in its entirety or in part submitted it for obtaining any qualification.

Signature:

Date: March 2016

Copyright © 2016 Stellenbosch University

All rights reserved

SUMMARY

Hormone replacement therapy (HRT) is used by post-menopausal women to alleviate symptoms associated with decreased endogenous estrogen levels, such as hot flashes and vaginal dryness. Although HRT is considered an effective treatment method, results from clinical trials such as those conducted by the Women's Health Initiative (WHI) and Million Women Study (MWS) revealed that HRT usage is associated with increased risk of invasive breast cancer. The mechanism whereby the hormones used in HRT contribute to breast cancer risk is not fully understood. These hormones elicit their biological effects by binding to intracellular steroid receptors such as the estrogen (ER) and progesterone receptor (PR). For many years it was thought that the ER is the main steroid receptor implicated in breast cancer biology, however, data in the literature suggests that crosstalk between steroid receptors play an important role in the development and progression of this disease. Recent evidence suggests that the androgen receptor (AR) can function as a tumour suppressor and thus has the potential of being a prognostic marker and therapeutic target in breast cancer. Given that the AR has been shown to inhibit the transactivation function of ER α , this raised the question of whether the AR would also inhibit the transactivation function of the ER β subtype. Moreover, considering that ERs have both transactivation and transrepression functions, studies investigating the influence of the AR on the transrepression function of both ER α and ER β are lacking. This study therefore investigated whether the AR, in the absence or presence of known AR agonists (the natural AR ligand 5 α -dihydrotestosterone (DHT)), the androgenic progestins, medroxyprogesterone acetate (MPA) and norethisterone-acetate (NET-A)), could modulate the transcriptional activity of the ER β . The ER- and AR-negative MDA-MB-231 breast cancer cell line, transiently transfected with expression vectors for either ER α or ER β and the appropriate promoter-reporter construct, was used as model system. The results showed that the unliganded or liganded AR has no effect on the transactivation function of ER β on a synthetic estrogen response element (ERE)-containing promoter but downregulates the ER β -driven mRNA expression of the endogenous PR gene. This study also shows estradiol (E₂)-induced transactivation on a synthetic ARE- and an endogenous androgen response element (ARE)-containing promoter via ER α . For transactivation via ER β , similar results were only seen on the endogenous ARE-containing promoter. Moreover, we show for the first time that both the unliganded- and ligand-bound AR inhibits the transrepression function of ER α , while interestingly only the ligand-bound AR was able to inhibit the transrepression function of ER β . No E₂-induced cell proliferation was observed in the MDA-MB-231 cell line overexpressing either ER α or ER β under the experimental conditions used. In conclusion, the results of this study are in agreement with previous evidence suggesting that the AR inhibits the activity of ER α . The precise physiological implications of these results remain to be determined. Finally, although the results from this study are preliminary and have certain limitations, it nonetheless highlights the fact that the role of the AR in breast cancer is a complex one. Thus, our findings may aid our understanding of crosstalk between the ER and AR signalling pathways, and how it contributes to the growth and survival of breast cancer cells.

OPSOMMING

Hormoon vervangingsterapie (HVT) word deur na-menopousale vrouens gebruik om simptome geassosieerd met verlaagde endogeen estrogeenvlakke, soos warmgloede en vaginale droogheid, te verlig. Alhoewel HVT beskou word as 'n effektiewe behandelingsmetode, toon resultate van kliniese toetse soos byvoorbeeld die Vroulike Gesondheids Inisiatief (VGI) en Miljoen Vroue Studie (MVS) dat die gebruik van HVT 'n toename in die risiko van indrigende borskanker veroorsaak. Die meganisme waardeur hierdie hormone, wat gebruik word in HVT, bydra tot die risiko van borskanker word nog nie ten volle verstaan nie. Hierdie hormone voer hul biologiese effekte uit deur te bind aan intrasellulêre steroïd reseptore soos die estrogeen (ER) en progesteron (PR) reseptore. Vir baie jare was dit geglo dat die ER die hoof steroïd reseptor betrokke is in borskanker, maar data in die literatuur stel egter voor dat 'n wisselwerking tussen steroïd reseptore 'n belangrike rol in die ontwikkeling en bevordering van hierdie siekte speel. Onlangse bewyse dui daarop dat die androgeen reseptor (AR) as 'n onderdrukker van gewasse kan funksioneer, en dus die potensiaal het om as 'n prognostiese merker en terapeutiese teiken in borskanker te dien. Die feit dat die AR al getoon het om die transaktiverings funksie van die ER α te onderdruk, het gelei tot die opwekking van die vraag of die AR ook die transaktiverings funksie van die ER β sub tipe kan onderdruk. Verder, siende dat die ER's beide transaktiverings en transonderdrukkings funksies het, is dit belangrik om in ag te neem dat daar 'n te kort is aan navorsing wat die invloed van die AR op die transonderdrukkings funksie van beide ER α en ER β ondersoek. Die huidige studie het dus 'n ondersoek ingestel om te bepaal of die AR, in die afwesigheid of teenwoordigheid van bekende AR agoniste (die natuurlike AR ligand 5 α -dihidrotestosteroon (DHT), die androgeniese progestiene medroksieprogesteron asetaat (MPA) en noretisteron asetaat (NET-A)), die transkripsionele aktiwiteit van die ER β kan verander. Die ER- en AR-negatiewe MDA-MB-231 borskankersellyn, tydelik getransfekteer met die uitdrukkingsvektore vir óf die ER α óf die ER β en die toepaslike promotor-rapporteerder konstrue, was gebruik as die model sisteem. Die resultate het getoon dat die AR geen effek op die transaktiverings funksie van die ER β op 'n sintetiese estrogeen respons element (ERE)-bevattende promotor het nie, maar dat dit wel die ER β -gedrewe mRNS uitdrukking van die endogene PR geen onderdruk. Estradiol (E₂)-geïnduseerde transaktivering op 'n sintetiese androgeen response element (ARE) en 'n endogene ARE-bevattende promotor deur ER α word ook in hierdie studie aangedui. Soortgelyke resultate is ook waargeneem vir die transaktivering deur ER β op 'n endogene ARE-bevattende promotor. Verder, wys ons vir die eerste keer dat beide die ligandlose en ligand-gebonde AR die transonderdrukkings funksie van ER α inhibeer, terwyl interessant genoeg slegs die ligand-gebonde AR die vermoë het om die transonderdrukkings funksie van ER β te inhibeer. Onder hierdie gebruikte eksperimentele kondisies, was geen E₂-geïnduseerde selproliferasie in die MDA-MB-231 sellyn, wat óf ER α óf ER β ooruitdruk, waargeneem nie. In samevatting, die resultate van hierdie studie stem ooreen met vorige bewyse wat voorstel dat die AR die aktiwiteit van ER α onderdruk. Die presiese fisiologiese implikasies van hierdie resultate moet nog bepaal word. Ten slotte, alhoewel die resultate van hierdie studie slegs voorlopig is en

sekere beperkinge toon, beklemtoon dit nietemin die feit dat die rol van die AR in borskanker kompleks is. Dus, ons bevindings mag bydra tot die huigelike begrip van die wisselwerking wat tussen die ER en die AR seintransduksiepaaie voorkom, en mag ook help om te verstaan hoe dit moontlik kan bydra tot die groei en oorlewing van borskankerselle.

I would like to dedicate this thesis to my dad, **Thamsanqa Joseph Ndlovu.**

May your soul rest in peace.

Philippians 4:8-9

ACKNOWLEDGEMENTS

I would like to greatly thank the following people who helped me to make this study possible:

First of all I would like to say a big thank you to my supervisor, Dr Donita Africander, for giving me this opportunity to do my masters under her supervision. She has been a great supervisor and I really adore and appreciate her remarkable dedication to her work. You are a true inspiration to many and you will definitely go far. It was really a great honour to go through this journey with you, although I know there were so many hurdles, you were always there to motivate me. I will always appreciate the way, you were always concerned and willing to listen during the toughest times. Words cannot express the way I really appreciate what you did for me.

Dr Renate Louw-Du Toit, thank you so much. You were always there for me since the first day I started this journey. Thank you for being the great teacher you were, no one could have done it better than you. The love and support you gave me was amazing. Thank you for being the sister you are and for all the times we used to talk you would motivate me to keep strong and push forward.

Mrs Carmen Langeveldt, thank you for all the hours you spent maintaining the cells in tissue culture. To me you are a hero, you always make sure that the lab is running smoothly.

I would also like to thank my co-supervisor, Professor Ann Louw for the help she gave me. Thank you for the advice and encouragement to keep moving forward, it was greatly appreciated.

Members of the Africander and Louw lab, thank you to Dr Nicolette Verhoog, Dr Koch Visser, Meghan Perkins, Legh Wilkinson, Morne Mortimer, Yolandi van der Meer, Craig Andrews, Herzelle Classen, and Kezia Klink.

My parents, my late dad, Joseph and mom Sibangani, I would like to thank you for all that you have done for. You made sure that I got a decent education. To my sisters my sisters Barbara, Sandisiwe and Manurse thank you for being there when I needed someone to talk.

I would also like to thank the following people, for the spiritual motivation they gave me, Pastor Funlola Olojede, Pastor Zachariah Bulus, Deacon John Babafemi and Sister Martha Bulus.

To the **LORD, GOD ALMIGHTY**, thank you for keeping me until this day. You are my pillar of strength.

LIST OF ABBREVIATIONS

17 β -HSD	17 β -hydroxysteroid dehydrogenase
AF-1	activation function-1
AF-2	activation function-2
AI	aromatase inhibitor
ANOVA	analysis of variance
AP-1	Activator Protein-1
AR	androgen receptor
ARA	androgen receptor associated
ARE(s)	androgen response element(s)
ATCC	American Type Culture Collection
AVG	average
bp	base-pair
CDK	cyclin dependent kinases
cDNA	complementary deoxyribonucleic acid
C/EBP	CCAAT enhancer binding protein
CHD	coronary heart disease
ChIP	chromatin immunoprecipitation
Co-IP	co-immunoprecipitation
CP	crossing point
CT	cycle number
C-terminal	carboxy-(COOH-) terminal
CTSD	cathepsin D

CYP	cytochrome P450
DBD	DNA-binding domain
DEPC	diethylpyrocarbonate
DHT	dihydrotestosterone
DMEM	Dulbecco's Modified Eagle Medium
DMSO	dimethyl sulfoxide
DNA	deoxyribonucleic acid
E	exponential amplification value
E ₁	estrone
E ₂	17 β -estradiol
E ₃	estriol
ECL	enhanced chemiluminescence
EDTA	ethylenediaminetetra-acetic acid
ER	estrogen receptor
ERE	estrogen response element
ER α	estrogen receptor alpha
ER β	estrogen receptor beta
EtOH	ethanol
FCS	fetal calf serum
FRET	fluorescence resonance energy transfer
GAPDH	glyceraldehyde 3-phosphate dehydrogenase
GF	growth factor
GnRH	gonadotropin-releasing hormone
GPCRs	G-protein-coupled receptors
GR	glucocorticoid receptor

HCL	hydrogen chloride
HER-2	human epidermal growth factor receptor-2
HRP	horseradish peroxide
HRT	hormone replacement therapy
Hsp(s)	heat-shock protein(s)
ICI 182,780	fulvestrant or faslodex
IκB	inhibitor of kappa B
IKK	IκB kinase
IL-6	interleukin-6
KLK	kallikrein
LB	Luria-Bertani
LBD	ligand binding domain
MAPK	mitogen-activated protein kinase
MIB	mibelorone
MNAR	modulator of non-genomic activity of estrogen receptor
MPA	medroxyprogesterone acetate
MR	mineralocorticoid receptor
mRNA	messenger ribonucleic acid
MTT	3-(4,5-dimethylthiazolyl-2)-2,5-diphenyltetrazolium bromide
MWS	Million Women Study
NET-A	norethisterone / norethindrone acetate
NFκB	nuclear factor kappa-B
NKX3	prostate homeobox
NO	nitric oxide
NRs	nuclear receptors

OD	optical density
PAGE	polyacrylamide gel electrophoresis
PBS	phosphate buffered saline
PMA	phorbol-myristate acetate
PR	progesterone receptor
PR-A	progesterone receptor A
PR-B	progesterone receptor B
PRE	progesterone response element
PSA	prostate specific antigen
qPCR	quantitative polymerase chain reaction
RANTES	Regulated on Activation, Normal T-cell Expressed, and Secreted
re-ChIP	chromatin re-immunoprecipitation
RLU	relative light unit
mRNA	messenger ribonucleic acid
RT-PCR	real time-polymerase chain reaction
SARM	selective androgen receptor modulator
SD	standard deviation
SDS	sodium dodecyl sulphate
SEM	standard error of the mean
SERD	selective estrogen receptor downregulator
SERM	selective estrogen receptor modulator
SHBG	sex hormone binding globulin
SM	second messenger
Sp1	specificity protein 1
SRC-1	steroid receptor co-activator 1

SRE	steroid response element
SRs	steroid receptors
R	relative expression
TBS	tris buffered saline
TBST	tris buffered saline tween
TFs	transcription factors
TGF- α	transforming growth factor alpha
T _m	melting temperature
TNBC	triple negative breast cancer
TNF- α	tumour necrosis factor alpha
VEGF	vascular endothelial growth factor
WHI	Women's Health Initiative
WHO	World Health Organization

TABLE OF CONTENTS

SUMMARY.....	iii
OPSOMMING.....	iv
ACKNOWLEDGEMENTS	vii
LIST OF ABBREVIATIONS	ix
TABLE OF CONTENTS	xiv
 Chapter 1 Literature Review	 1
1.1. Introduction.....	2
1.2. Molecular mechanism of action of steroid receptors	4
1.2.1. General structure of steroid receptors	4
1.2.2. Transactivation.....	7
1.2.3. Transrepression	7
1.3. Molecular mechanisms of estrogen action.....	9
1.3.1. Production of estrogens and their physiological actions.....	9
1.3.2. Diverse molecular pathways are involved in ER-mediated estrogen actions	13
1.4. The role of E₂ and ER subtypes in breast cancer biology.....	17
1.4.1. E ₂ and ER α	17

1.4.2. ER β and its interplay with ER α	20
1.5. Role of androgens and the AR in breast cancer biology	21
1.5.1. Androgens	21
1.5.2. Androgen receptor (AR)	23
1.5.3. Crosstalk between the ER subtypes and the AR.....	25
1.6. Conclusion	28
1.7. Aims of the study.....	29
 Chapter 2 Materials and Methods	 31
2.1. Plasmids	32
2.1.1. Plasmid DNA preparation.....	32
2.2. Test compounds.....	33
2.3. Cell culture	34
2.4. Promoter reporter assays.....	34
2.4.1. Transactivation.....	34
2.4.2. Transrepression	36
2.5. Western blot analysis.....	36
2.6. Total RNA isolation and complementary DNA (cDNA) synthesis...38	
2.7. Realtime quantitative polymerase chain reaction (qPCR)	40

2.8. Methyl Thiazolyl Tetrazolium (MTT) cell proliferation assay	42
2.9. Data manipulation and statistical analysis	42
Chapter 3 Results	44
3.1. Validating the experimental system	45
3.2. The AR does not affect the transcriptional activation of ER β on a synthetic ERE-containing promoter	48
3.3. The AR ligands do not augment the effect of the unliganded AR ...	51
3.4. AR abolishes the transcriptional activity of E ₂ via both ER subtypes on an endogenous ER regulated gene in a breast cancer cell line....	57
3.5. Androgen response element transactivation via ER α but not ER β .	61
3.6. E ₂ -induced upregulation of an endogenous ARE-containing gene can be mediated by both ER α and β	63
3.7. The AR inhibits the transrepression function of both ER α and ER β	65
3.8. No change in proliferation was observed in cells transfected with ER	73
Chapter 4 Discussion	75

4.1. Introduction.....	76
4.2. Validating the model system	77
4.3. Effects on synthetic promoter-reporter genes are not always reflected on endogenous genes.....	78
4.4. E₂ upregulation of the endogenous ARE-containing PSA gene can be mediated by either ERα or ERβ	80
4.5. The unliganded AR impedes repression of transcription via ERα, but not ERβ, while the liganded AR impedes the transrepression function of both ERα and ERβ.	82
4.6. No change in proliferation was observed in cells transfected with ER	83
4.7. Conclusions and Future Work	84
 REFERENCES	 90
 ADDENDUM	 130

Chapter 1

Literature Review

1.1. Introduction

Breast cancer accounts for about one fourth of all female cancers, and is the second leading cause of cancer-related death worldwide in women (Sommer and Fuqua 2001, Wang *et al.* 2006, Yu *et al.* 2011, Seok-Jun *et al.* 2012, Chottanapund *et al.* 2013). Several factors have been identified, which increases the risk of developing breast cancer, and include genetics, epigenetics, lifestyle and endocrine therapy such as hormone replacement therapy (HRT) (Bergkvist *et al.* 1989, Black 1994, Colditz *et al.* 1995, McPherson *et al.* 2000, Rossouw *et al.* 2002, Beral 2003). HRT is used by post-menopausal women to alleviate the symptoms associated with menopause, due to decreased estrogen levels. The symptoms of menopause include hot flashes, vaginal dryness, dysfunctional uterine bleeding, urogenital atrophy and psychological symptoms such as irritability, depression and insomnia (McKinlay *et al.* 1992, Ravn *et al.* 1994, Farquhar *et al.* 2009). HRT is given as estrogen alone to women who have undergone a hysterectomy, while women with an intact uterus are administered a combination of estrogen and a progestin, such as medroxyprogesterone acetate (MPA) or norethisterone acetate (NET-A) (Pike *et al.* 1997, Marsden 2003, Verkooijena *et al.* 2009, Mia *et al.* 2013). Progestins are added to reduce the risk of estrogen-induced endometrial hyperplasia (Gelfand and Ferenczy 1989, Pinto *et al.* 2003, Furness *et al.* 2009). Although HRT is considered an effective treatment for relieving menopausal symptoms, results from the Women's Health Initiative (WHI) trial and the 'Million Women Study' (MWS) raised much alarm by showing that the use of HRT (estrogen alone and estrogen plus progestin) leads to increased risk of breast cancer in women (Rossouw *et al.* 2002, Beral 2003). Interestingly, recent studies have shown that there has been a decrease in breast cancer incidence, which has been suggested to be linked to a decline in HRT usage (Katalinic and Rawal 2008, Silvermana *et al.* 2011, Antoine *et al.* 2014). It is thought that the primary cause of the reduction in the use of HRT over the last few years is due to the results of the WHI study (Chlebowski *et al.* 2003, Chlebowski *et al.* 2010, Holm *et al.* 2014). Understanding the mechanisms whereby hormones used in HRT increase

women's risk of developing breast cancer is essential for finding improved therapies for the treatment of breast cancer. It is known that the hormones used in HRT elicit their biological effects primarily via binding to the estrogen receptor (ER) or the progesterone receptor (PR), which are both members of the steroid receptor family (Osborne *et al.* 2001, Zilli *et al.* 2009, Kittler *et al.* 2013, Knutson and Lange 2014). The ER exists as two subtypes namely ER α and ER β (Brandenberger *et al.* 1997, Enmark *et al.* 1997, Nilsson *et al.* 2001), while the PR is expressed as two isoforms called PR-A and PR-B, respectively (Kastner *et al.* 1990).

Estrogen acting via ER α is considered to be the major etiological factor in breast cancer and stimulates breast cancer cell proliferation in ER-positive breast cancer cells (Jordan and Ford 2011, Wu *et al.* 2012), while ER β inhibits ER α -mediated cell proliferation (Gougelet *et al.* 2005, Paris *et al.* 2012, Saxena *et al.* 2013). Studies have reported that the PR is also involved in breast cancer cell proliferation (Kalkhoven *et al.* 1994, Saitoh *et al.* 2005, Chen *et al.* 2012), with differential contributions of the PR isoforms being reported (Graham *et al.* 1996, Hopp *et al.* 2004, Stendahl *et al.* 2006, Giulianelli *et al.* 2013). For example, PR-B has been shown to have proliferative effects on breast cancer cell proliferation in the presence of its natural ligand progesterone (Conneely *et al.* 2003, Mulac-Jericevic *et al.* 2003, Boonyaratanakornkit *et al.* 2008, Lanari *et al.* 2012), while PR-A has anti-proliferative properties (Saitoh *et al.* 2005, Boonyaratanakornkit *et al.* 2008, Khan *et al.* 2012). Although the ER and the PR are the best-known prognostic markers of breast cancer (Silvermana *et al.* 2011), other steroid receptors such as the androgen receptor (AR) as well as the glucocorticoid receptor (GR), have been implicated in the etiology and pathogenesis of hormone-dependent breast cancer (Teulings *et al.* 1980, Conzen 2008). In fact, current evidence in the literature suggests that both the AR and GR display anti-proliferative effects and may thus have protective roles in breast cancer (Isola 1993, Kuenen-Boumeester *et al.* 1996, Moinfar *et al.* 2003, Katalinic and Rawal 2008, Peters *et al.* 2009, Robinson *et al.* 2011).

It is noteworthy that some steroid receptors have been reported to be co-expressed in breast cancer tumours (Teulings *et al.* 1980, Ciocca *et al.* 2006, Gago *et al.* 2006, McNamara *et al.* 2013). For example, it is known that the AR can be co-expressed with the ER in breast cancer tumours (Kuenen-Boumeester *et al.* 1996, Moinfar *et al.* 2003, Castellano *et al.* 2010). Interestingly, the expression of the AR in ER-positive breast cancer tumours is associated with a more favourable prognosis (Moinfar *et al.* 2003, Alshenawy 2012, Qu *et al.* 2013). Since the AR is expressed in most tumours, and as current therapies targetting estrogen signalling pathways have been shown to be ineffective in some patients due to the development of drug resistance (Martin *et al.* 2005, Ciruelos *et al.* 2014), a drug targetting the AR signalling pathway, could potentially be beneficial. It is important to understand the AR signalling pathways so that they can be targetted for the development of improved breast cancer therapies (Brentani *et al.* 1986, Birrell *et al.* 1998, McNamara *et al.* 2013). Therefore the objective of this literature review is to highlight the respective roles of the ER subtypes and the AR in breast cancer biology, by focusing on the mechanism of action of these steroid receptors when bound to either cognate or synthetic ligands. The implications of crosstalk between these receptors for breast cancer biology will also be discussed.

1.2. Molecular mechanism of action of steroid receptors

The intracellular effects of steroid hormones are mediated via binding to members of the steroid receptor family, a subfamily of the nuclear receptor family, and includes the ER, PR, AR, GR and mineralocorticoid receptor (MR) (Gasc and Baulieu 1987, Couse and Korach 1999, Lu *et al.* 2006, Helsen and Frank 2014). These receptors are ligand-activated transcription factors with similarities in their structure and function.

1.2.1. General structure of steroid receptors

As shown in figure 1.1, steroid receptors consist of five distinct functional domains, designated A-E, with ER α having an additional, carboxy-terminal F domain (Sommer and Fuqua 2001,

Balfe *et al.* 2004, Yang *et al.* 2008, Charitidi *et al.* 2009, Jin and Li 2010). The amino-terminal A/B domain contains the variable ligand-independent activation function (AF)-1 domain, which is important for protein-protein interaction with transcription factors and cofactors. This is followed by a highly conserved and centrally located DNA binding domain (DBD) (C-domain), a hinge region (D-domain) and a moderately conserved ligand binding domain (LBD) (E-domain) situated at the carboxy-terminal (figure 1.1) (Jin and Li 2010, Forrest and Swaroop 2012); reviewed in (Bain *et al.* 2007); reviewed in (Africander *et al.* 2011)).

The DBD is highly homologous between the steroid receptors and comprises two zinc-finger motifs. The α -helix in the first zinc finger plays an important role in enabling the DBD to recognize specific palindromic DNA sequences known as steroid response elements (SREs) (Saji *et al.* 2005, Wu and Safe 2007, Charitidi *et al.* 2009, Helsen *et al.* 2012, McEwan 2012), while the second zinc finger is important for receptor homo- or heterodimerization (Shaffer and Gewirth 2002, Helsen and Frank 2014). The DBD is linked to the LBD via a short highly variable hinge region (D domain) (Wu and Safe 2007, Helsen *et al.* 2012, Hudson *et al.* 2014), which contains sites for post-translational modifications such as phosphorylation, methylation and sumoylation (Anbalagan *et al.* 2012, Helsen and Frank 2014). A second activation function domain (AF-2), is located in the LBD, and is necessary for recruiting various cofactors and for the induction of the transcriptional activity of the receptor (Kumar and Chambon 1988, Xu and Li 2003, Bain *et al.* 2007). It should be noted that the AF-2 transcriptional activity is ligand-dependent, while that of AF-1 is ligand-independent. Ligand-dependent recruitment of coregulators to the AF-2 domain occurs via a hydrophobic cleft in this domain, which is shielded by an amphipathic α -helix in the absence of ligand (Griekspoor *et al.* 2007). The presence of ligand repositions this helix, thereby allowing coregulator binding via conserved LXXLL motifs found in the AF-2 domain (Weatherman *et al.* 1999). The sequence homology among the LBDs of steroid receptors is moderately conserved, with about a 50-70% sequence homology between the AR, GR, MR and PR, while the LBDs of the ER and PR share only

about 28% sequence homology, and that of the GR and ER only about 30% (Fuller 1991, Moras and Gronemeyer 1998, Maeda 2001). Lastly, the F domain, unique to ER α , contains functional elements which are responsible for functions such as controlling the interaction of ER α with steroid receptor coactivator-1 (SRC-1), and plays an important role in the receptor's ability to distinguish between agonists and antagonists (Montano *et al.* 1995, Koide *et al.* 2007).




Figure 1.1. A schematic illustration of the general structure of steroid receptors. The A/B region denotes the variable N-terminal domain containing the activation function-1 (AF-1) transactivation domain, C is the conserved DNA binding domain (DBD), D denotes the hinge region, and E is the carboxy-terminal moderately conserved ligand binding domain (LBD) containing the AF-2 transactivation region. ER α , contains an additional F domain. The numbers represent the amino acid length of the steroid receptors. Figure taken from (Africander *et al.* 2011).

Most unliganded steroid receptors are found in the cytoplasm, however, the ER and PR are found mainly in the nucleus (Brosens *et al.* 2004). These unliganded steroid receptors are sequestered in an inactive form in a large molecular complex organised around the molecular chaperone heat shock proteins, Hsp90 and Hsp70 (Pratt and Toft 2003, Gougelet *et al.* 2007).

When activated by ligand, steroid receptors modulate gene expression via two mechanisms, namely transactivation and transrepression.

1.2.2. Transactivation

Ligand-dependent transcriptional activation of genes occurs when small lipophilic hormones diffuse across the plasma membrane and bind to the LBD of the steroid receptor, leading to a conformational change in the steroid receptor and translocation to the nucleus of those steroid receptors found predominantly in the cytoplasm (figure 1.2) (Zhao *et al.* 2010, Murphy 2011). Once in the nucleus, the hormone-bound steroid receptor binds as a dimer to specific SREs found in the promoters of steroid responsive target genes (Beato and Klug 2000, Kumar and Litwack 2009). For example, the ER binds to SREs called estrogen response elements (EREs), while the PR binds to progesterone response elements (PREs). The SREs for PR, AR, MR and GR are highly homologous and consist of the palindromic sequence 5'-GGTACAnnnTGTTCT-3' (Beato *et al.* 1989), while the ER recognizes the alternative 5'-GGTCAnnnTGACC-3' palindromic sequence (Klock *et al.* 1987, Klein-Hitpass *et al.* 1988, Brosens *et al.* 2004). The binding of a steroid receptor to the SRE, is followed by recruitment of coregulators, chromatin remodelling, leading to the recruitment of the basal transcriptional machinery (Beato and Klug 2000, Perissi and Rosenfeld 2005, Lu *et al.* 2006). This leads to the activation of gene transcription in a process known as transactivation (figure 1.2).

1.2.3. Transrepression

Steroid receptors can also negatively regulate transcription of genes by binding to DNA-bound transcription factors such as nuclear factor kappa-B (NFκB), activator protein-1 (AP-1) and CCAAT enhancer binding protein (C/EBP) (Ueda *et al.* 1994, Harrington *et al.* 2003, Pascual and Glass 2006, Frasor *et al.* 2009) in a process called transrepression (Harrington *et al.* 2003,

Lu *et al.* 2006, Carter and Stephen 2014). Although the ER, AR, PR and MR have all been shown to interact with NF κ B to repress the expression of target genes (Kalkhoven *et al.* 1996, Palvimo *et al.* 1996, Ray *et al.* 1997, Aarnisalo *et al.* 1998, McKay and Cidlowski 1999, Kalaitzidis and Gilmore 2005, S nder *et al.* 2006, Matter and Handschin 2007, Bergmann *et al.* 2010, Chantong *et al.* 2012, Drummond and Fuller 2012, Africander *et al.* 2014), the mechanism of transrepression is best studied for the GR through its interaction with NF κ B and AP-1 (McKay and Cidlowski. 2000, Pascual and Glass 2006, Newton and Holden 2007, De Bosscher *et al.* 2008, Beck *et al.* 2009). NF κ B is a transcription factor that is activated by many stimuli such as the tumor promoter, phorbol 12-myristate 13-acetate (PMA) (Lee *et al.* 2002, Chang *et al.* 2005) and proinflammatory cytokines such as tumour necrosis factor- α TNF- α (Scarpati and Sadler 1989, Ravi and Bedi 2004, Lee *et al.* 2007, Beck *et al.* 2009). Upon ligand binding to the GR, the liganded GR translocates to the nucleus where it tethers to DNA-bound NF κ B, thereby inhibiting NF κ B-driven gene expression (figure 1.2) (Ghosh *et al.* 1998, De Bosscher *et al.* 2008). Similar to the mechanism of transactivation, but less understood, transrepression also involves the binding of multiprotein complexes on DNA and the modulation of chromatin structure (Kassel and Herrlich 2007, Africander *et al.* 2011). Considering that all steroid receptors have essentially identical mechanisms of action for transactivation, it is probable that they repress genes by a similar mechanism to that of the GR. As this study focuses on the transcriptional activity of the ER and AR, a more detailed discussion on these receptors will follow.




Figure 1.2. An illustration of the classical (general) mechanism of action of steroid receptors. SR: steroid receptor, SRE: steroid response element, Hsp90: heat shock protein 90, NFκB: nuclear factor kappa B, AP-1: activator protein. Co-regulators (represented by cofactors in the diagram), result in the opening of the chromatin structure and recruitment of basal transcription machinery. Redrawn from (Africander *et al.* 2011).

1.3. Molecular mechanisms of estrogen action

1.3.1. Production of estrogens and their physiological actions

Estrogens are female sex steroid hormones that play an important role in various tissues of the human body (Gougelet *et al.* 2005, Thomas and Potter 2013). Figure 1.3 shows the many examples of these roles, which include maintenance of bone density as well as the growth and proliferation of breast tissue (Gruber *et al.* 2002).




Figure 1.3. Estrogens act on multiple organ systems. The figure illustrates the diverse actions of estrogens on different organs in the human body. Figure taken from (Gruber *et al.* 2002).

Three types of estrogens are produced within the female body namely, estrone (E_1), 17 β -estradiol (E_2), and estriol (E_3) (Thomas and Potter 2013, Samavat and Kurzer 2015). E_2 is mainly produced in the ovaries (Simpson 2002, Tsuchiya *et al.* 2005), while E_1 and E_3 are primarily produced in the liver from E_2 (Gruber *et al.* 2002, Tsuchiya *et al.* 2005). As shown in figure 1.4, E_2 is formed from testosterone either directly by the action of the aromatase (CYP19) enzyme, or indirectly through the reduction of E_1 by 17 β -hydroxysteroid dehydrogenase (17 β -HSD) isozymes. The 17 β -HSD2 enzyme oxidises testosterone to androstenedione, which is

aromatized to E₁ (Tsuchiya *et al.* 2005, Knower *et al.* 2013, Thomas and Potter 2013, Samavat and Kurzer 2015). E₃ is synthesized via the action of the enzyme 17 β -HSD1 on the intermediate 16 α -hydroxyestrone, which is produced from E₁ via cytochrome P450 3A4/5 (CYP3A4 or CYP3A5) (Huang *et al.* 1998). E₁ is the inactive form of E₂ (Andersson and Moghrabi 1997, Abplanalp *et al.* 1999) and has been shown to inhibit cell division of the MCF-7 breast cancer cell line (Jozan *et al.* 1979). Although previous studies have suggested that high levels of E₃ protect women against breast cancer (Lemon 1975, Wotiz *et al.* 1978, Lemon *et al.* 1989), some studies have shown that E₃ can increase the growth of breast cancer cells and activate expression of ERE-containing genes to the same extent as E₂ (Katzenellenbogen 1984, Pang and Faber 2001, Diller *et al.* 2014). As E₂ is the most biologically active estrogen, with circulating levels in the high picomolar to nanomolar range in the reproductive phase of women (Chatterton *et al.* 2003, Watson *et al.* 2008, Thomas and Potter 2013, Samavat and Kurzer 2015), we will only focus on the actions of E₂.




Figure 1.4 legend on next page.

Figure 1.4. An illustration of the estrogen metabolism pathway. Testosterone is the main precursor for estrogen production, which involves different cytochrome P450 enzymes. Aromatase is the major enzyme involved in the production of E₂ and E₁ from the androgen precursors (testosterone and androstenedione), 17 β -HSD plays a role in the production of E₁ and E₃, while CYP3A4/5 is involved in the production of E₃ from E₁. Adapted from (Thomas and Potter 2013).

E₂ is released into the bloodstream after production and ~37% binds to sex hormone binding globulin (SHBG) and ~60% binds to albumin for transport to target tissues (Dunn *et al.* 1981), while only ~3% is left free or unbound (Siiteri 1987, Pardridge 1988, Petra 1991, Gruber *et al.* 2002). Free E₂ diffuses into target tissues where its biological actions are primarily mediated by binding to estrogen receptors (Matthews and Gustafsson 2003, Balfe *et al.* 2004, Faulds *et al.* 2012), and these actions include the normal female reproductive processes as well as playing a role in the regulation of a number of diseases. Examples of the latter include osteoporosis, coronary heart disease (CHD), Alzheimer's disease and hormone-dependent cancers, which include endometrial, ovarian, prostate and breast cancer (Ingemar 2000, Paranjape *et al.* 2005, Li-Min *et al.* 2012, Wen-Yang *et al.* 2012, Brown and Hankinson 2014, Li *et al.* 2014, Hamed and Mindy 2015). It is important to note that the role of E₂ in the above-mentioned processes can be either positive or negative. Examples of positive effects include reports that high levels of E₂ prevents the development of Alzheimer's disease (Giacobini 1998, Xing *et al.* 2013, Li *et al.* 2014), while the use of E₂ plus progestin in HRT has been associated with a decreased risk of osteoporosis (Lee *et al.* 2013, Nelson *et al.* 2013) and coronary heart disease (CHD) (Grodstein *et al.* 1996, Paranjape *et al.* 2005, Hodis and Mack 2014). In contrast to these positive effects, E₂ has been shown to increase the risk of endometrial and ovarian cancer via several mechanisms such as cell proliferation, decreased apoptosis and increased DNA damage (Henderson and Feigelson 2000, Cunat *et al.* 2004, Liang *et al.* 2009, Zhang *et al.* 2012, Brown and Hankinson 2014, Ribeiro and Freiman 2014). Similarly, long-term exposure to E₂ has been shown to be linked to an increased risk of breast cancer and may lead to tumor growth via

mechanisms such as increased DNA damage and cell proliferation (Fox *et al.* 2008, Santen *et al.* 2009, Yue *et al.* 2013).

1.3.2. Diverse molecular pathways are involved in ER-mediated estrogen actions

1.3.2.1. ER subtypes

The main actions of estrogen are genomic and are mediated by the ER subtypes. The two main functional ER subtypes, ER α and ER β , are transcribed from different genes (Menasce *et al.* 1993, Matthews and Gustafsson 2003). ER α is a 66 kDa protein, which was discovered in the late 1960s (Toft *et al.* 1967) and was isolated and purified from the human MCF-7 breast cancer cell line in the late 1980s (Walter *et al.* 1985, Green *et al.* 1986). It is expressed in the female reproductive system (breast, ovary, uterus), male reproductive system (testis, epididymis, prostate gland) and various other tissues such as the pituitary gland, kidney, bladder, liver, heart, thymus and adrenal gland (Ogawa *et al.* 1998, Pearce and Jordan 2004). ER β is a 56 kDa protein, which was only identified in 1996 (Kuiper *et al.* 1996, Mosselman *et al.* 1996, Katzenellenbogen and Korach 1997, Enmark and Gustafsson 1999), and like ER α , ER β is expressed in the female and male reproductive systems as well as various other tissues such as the bladder, lung, adrenal and heart (Iwao *et al.* 2000, Roger *et al.* 2001, Farquhar *et al.* 2009).

Human ER α and ER β have similar structural domains (figure 1.5) and differ mainly in their amino acid length (Nilsson *et al.* 2001, Pearce and Jordan 2004, Ascenzi *et al.* 2006). The N-terminal domain of the ERs has only about 30% identity, with ER β being reported to lack AF-1 activity (Pearce and Jordan 2004). The DBD of the two ERs has 96% identity, suggesting that both receptors would bind to similar sites in target genes. The hinge region, like the N-terminal domain, is not highly conserved between the ERs (30% homology). The LBD's of the two subtypes have a 53% homology, and show slight differences in ligand binding specificity

(Mosselman *et al.* 1996, Pearce and Jordan 2004). For example, E₂ has been shown to have a slightly higher binding affinity for ER α as compared to ER β , while phytoestrogens such as genistein has a higher binding affinity for ER β (Verhoog *et al.* 2007).




Figure 1.5. Structural homology between human ER α and ER β . The various domains are represented by the letters A-F, with the amino acid sequence homology between the individual domains shown as a percentage. The A/B region denotes the variable N-terminal domain containing the AF-1 domain, C is the DBD, D denotes the hinge region which connects the DBD to the LBD, E is the LBD, which contains the AF-2 domain. ER α , contains an additional F domain. The numbers represent the amino acid length of the steroid receptors. Figure redrawn from (Pearce and Jordan 2004, Africander *et al.* 2011).

1.3.2.2. Molecular pathways

ERs can regulate biological processes through several distinct signaling pathways, which include both ligand-dependent and ligand-independent mechanisms (figure 1.6). The classical genomic mechanism of steroid receptor action is ligand-dependent (illustrated in figure 1.2), and for the ER, involves E₂ binding to the ER, and the subsequent direct binding of the E₂-bound ER to an ERE (figure 1.6A) or tethering to another transcription factor (figure 1.6B) (Gruber *et al.* 2002, Simoncini *et al.* 2004, Yager and Davidson 2006, Heldring *et al.* 2007).

These nuclear actions lead to increased gene expression of target genes via the ERE, and

decreased transcription of target genes via tethering to DNA-bound transcription factors such as NF κ B and AP-1 (Stein and Yang 1995, Galien and Garcia 1997, Yamei *et al.* 2007, Robertson *et al.* 2010). Interestingly, the ER has been shown to repress NF κ B signalling at various levels of transcriptional regulation (Kalaitzidis and Gilmore 2005, Drummond and Fuller 2012). For example, the ER can repress NF κ B signaling by (i) suppressing inhibitor of kappa B (IkB) kinase (IKK) activity, (ii) preventing the degradation of IkB proteins, (iii) blocking NF κ B from binding to DNA and (iv) competing with NF κ B for coactivators (Kalaitzidis and Gilmore 2005, Drummond and Fuller 2012). However, there is limited information in the literature regarding ER subtype-specific-mediated mechanisms of transrepression. At least one study has shown that E₂ acting via both ER subtypes, can repress the expression of an NF κ B containing promoter-reporter construct (Visser *et al.* 2013), while one other study showed that E₂ acting via ER α was able to cause repression of three TNF- α -induced proinflammatory genes (Cvoro *et al.* 2011).

Although the classical mechanism of action can take hours to achieve maximal E₂-induced gene responses (Gruber *et al.* 2002, Simoncini *et al.* 2004, Heldring *et al.* 2007), some of the ligand-activated ER responses are more rapid (Watsona *et al.* 1999, Gruber *et al.* 2002, Wong *et al.* 2002, Simoncini *et al.* 2004, Song *et al.* 2005, Song and Santen 2006). An example of the latter is the non-genomic mechanism involving membrane-bound ER (figure 1.6C) and most likely involves other membrane bound receptors such as G-protein-coupled receptors (GPCRs). In this mechanism the ligand activates the membrane-associated ER, leading to the initiation of signaling cascades via second messengers (SM). These SM affect ion channels or increase nitric oxide (NO) levels in the cytoplasm, thereby causing rapid physiological responses without involving gene regulation (Hall *et al.* 2001, Stefan *et al.* 2001, Björnström and Sjöberg 2005, Heldring *et al.* 2007, Silva *et al.* 2010, Ozgyin *et al.* 2015). Lastly, growth factor (GF) signalling (figure 1.6D) can lead to the activation of protein kinase cascades, which result in the

phosphorylation of ERs and the subsequent binding to EREs, thus modulating gene regulation via a ligand-independent mechanism (Björnström and Sjöberg 2005, Heldring *et al.* 2007).




Figure 1.6. Molecular pathways used in the regulatory actions of ERs. The classical (direct) pathway (A) involves ligand-bound ER binding directly to EREs leading to increased transcription of target genes. Alternatively, the tethered pathway (B) involves indirect binding to DNA via protein-protein interactions with other transcription factors resulting in decreased transcription of target genes. The non-genomic pathway (C) involves ligand activation of membrane receptors, which leads to the initiation of signaling cascades via second messengers (SM), increasing nitric oxide (NO) levels and ultimately leading to altered physiological responses. Lastly the ligand-independent mechanism (D) involves activation via growth factor (GF) signalling leading to the activation of kinases that phosphorylate the ERs thereby activating them and modulating gene transcription. Figure modified from (Heldring *et al.* 2007).

1.4. The role of E₂ and ER subtypes in breast cancer biology

1.4.1. E₂ and ER α

Numerous clinical and epidemiological studies have indicated that exposure to E₂ is associated with increased risk of developing breast cancer (Clemons and Goss 2001, Yager and Davidson 2006). The cellular mechanisms by which E₂ increases breast cancer risk is an area of ongoing study but at least two possibilities have been suggested (figure 1.7) (Miller 2004). Firstly, E₂ binds to the ER causing altered gene expression, which results in uncontrolled or excessive cell proliferation (figure 1.7A). Interestingly, the E₂-induced proliferative effects on breast cancer cells in humans can be either direct or indirect (Clemons and Goss 2001). The direct E₂-induced mechanism of cell proliferation is initiated by the induction of enzymes and proteins involved in nucleic acid synthesis and activation of specific oncogenes (Clemons and Goss 2001), while the secretion of hormones such as prolactin and the production of GFs such as transforming growth factor alpha (TGF α) initiates the indirect mechanism of cell proliferation (Clemons and Goss 2001). The second mechanism whereby it has been suggested that E₂ causes increased risk of breast cancer comes from evidence suggesting that the metabolism of E₂ to genotoxic agents causes direct DNA mutations (figure 1.7B) (Yager and Leihhr 1996, Miller 2004, Yager and Davidson 2006). E₂ is metabolized via catechols (organic compounds), particularly 2- or 4- hydroxyestradiol or hydroxyestrone, to produce quinones. The latter interacts with and mutates DNA, initiating carcinogenesis (figure 1.7B) (Miller 2004, Yager and Davidson 2006). Considering that this thesis focusses on some parts of the pathway shown in figure 1.7A, and that the ER α subtype has predominantly been implicated in the development and progression of breast cancer, only the role of E₂ via the ER α will be further discussed.




Figure 1.7. Two pathways indicating the mechanisms of E₂ carcinogenesis. The diagram illustrates two mechanisms that contribute to the initiation of breast cancer. (A) ER-mediated increase in cell proliferation and (B) metabolism of E₂ to genotoxic agents such as quinones, causing DNA mutation. Redrawn from (Miller 2004).

ER α is expressed in almost 70% of breast cancers (Jensen and Jordan 2003, Pearce and Jordan 2004, Conzen 2008). Both *in vivo* and *in vitro* studies have reported that ER α is crucial for the promotion of breast cancer cell growth and survival by E₂ (Cunha *et al.* 1997, Couse and Korach 1999, Frasor *et al.* 2003, Hartman *et al.* 2009, Briskin and O'Malley 2010, Powell *et al.* 2012, Huang *et al.* 2014b). For example, despite the presence of high levels of E₂, no breast cancer tumours were detected in ER α knock-out mice (Couse and Korach 1999). Current therapies for E₂-dependent breast cancer are thus directed towards either (i) preventing the interaction of E₂ with the ER α using selective ER modulators (SERMs) or selective estrogen receptor downregulators (SERDs) (Jordan 2007, Patani and Martin 2014) or (ii) by inhibiting the synthesis of E₂ using aromatase inhibitors (AI's) (Mirzaie *et al.* 2013, Nantasenamat *et al.* 2013,

Akçay and Bayrak 2014). The latter includes compounds such as letrozole and anastrozole, which bind reversibly to the aromatase enzyme, thereby decreasing the levels of E₂ available to promote tumour formation or progression (Kijima *et al.* 2005, Gil *et al.* 2013). Studies have shown that anastrozole decreases cell proliferation in breast cancer cell lines (Miller and Jackson 2003, Kijima *et al.* 2005, Gil *et al.* 2013) and reduces tumour size when administered to breast cancer patients (Miller and Jackson 2003, Kijima *et al.* 2005, Gil *et al.* 2013). SERMs on the other hand are ligands that bind the ER α and act as either agonists or antagonists depending on the type of tissue (Howell *et al.* 2004, Renoir 2012). In the breast however, SERMs act as ER α antagonists (Macgregor and Jordan 1998, O'Regan and Jordan 2002) and include tamoxifen, the most frequently used drug in first-line clinical therapy for the treatment and prevention of hormone-sensitive breast cancer (Ogawa *et al.* 1998, Girault *et al.* 2006, Zervoudis *et al.* 2014, Lim *et al.* 2015), as well as raloxifene (Kijima *et al.* 2005, Zhang *et al.* 2013). Both these ligands have been shown to inhibit breast cancer cell growth (Speirs *et al.* 1999, Fowler *et al.* 2004, Howell *et al.* 2004, Speirs *et al.* 2004, Kijima *et al.* 2005, Lange 2008) by antagonizing E₂-induced gene expression and proliferation. SERDs are pure antagonists that do not have any agonist activity via the ER α in any tissue (Macgregor and Jordan 1998, O'Regan and Jordan 2002, Howell *et al.* 2004, Renoir 2012), and ICI 182,780, also known as fulvestrant or faslodex, and RU58668 (Pearce and Jordan 2004, Renoir 2012, Ciruelos *et al.* 2014). Clinically, faslodex is used as second-line treatment when patients fail to respond to tamoxifen (Lynn 2004, Martin *et al.* 2005, Howell 2006, Ciruelos *et al.* 2014). SERDs inhibit ER α -mediated transcriptional activity by down-regulating ER α levels, as these ligands target the ER α for proteasome-dependent degradation (Alarid *et al.* 1999, Lonard *et al.* 2000, Meiyun *et al.* 2003, Howell 2006, Renoir 2012, Ciruelos *et al.* 2014).

Several studies have shown that E₂-bound ER α promotes the development and progression of breast cancer by upregulating the expression of genes, such as c-myc, c-fos and cyclin D1, that play a role in the cell cycle and induce breast cancer cell (Helguero *et al.* 2005, Hartman *et al.*

2009, Briskin and O'Malley 2010), and/or by eliciting anti-apoptotic effects through the upregulation of the anti-apoptotic protein Bcl-x and downregulation of the pro-apoptotic protein Bax (Helguero *et al.* 2005). Not only does E₂ stimulate breast cell proliferation and inhibit apoptosis via ER α , but it also plays a role in cell invasion and metastasis (Clarke *et al.* 2003, Jiang *et al.* 2007). For example, E₂ stimulates the invasive and metastatic potential of the ER α -positive MCF-7 breast cancer cell line (Shafie and Liotta 1980, Osborne *et al.* 1985, Cos *et al.* 1998).

1.4.2. ER β and its interplay with ER α

Immunohistochemical analysis has shown that both ER α and ER β are expressed in the normal breast, and that ER β expression is more abundant than ER α (Clarke *et al.* 1997, Russo *et al.* 1999, Speirs *et al.* 2004). However, more ER α than ER β is expressed during the development of breast cancer (Leygue *et al.* 1998, Roger *et al.* 2001, Murphy *et al.* 2003, Shaaban *et al.* 2003). Interestingly, a small number of breast cancers express only ER α (~14%) or only ER β (~18%) (Murphy *et al.* 2003, Skliris *et al.* 2008, Leung *et al.* 2012). It is important to note that the current evidence in the literature suggests that co-expression of ER α and ER β in breast cancer patients is linked to a better prognosis than when ER α or ER β alone is expressed (Omoto *et al.* 2001, Roger *et al.* 2001, Paruthiyil *et al.* 2004, Lin *et al.* 2007). This is due to the ability of ER β to inhibit ER α -mediated cell proliferation as well as its ability to induce apoptosis (Leygue *et al.* 1998, Dotzlaw *et al.* 1999, Speirs *et al.* 1999, Zhao *et al.* 2008). For example, evidence from *in vitro* studies indicate that co-expression of ER β in ER α -positive breast cancer cells inhibits the proliferative effects of E₂ via ER α (Williams *et al.* 2007, Hartman *et al.* 2009). In fact, even in the absence of E₂, co-expression of ER β with ER α in the same cell line, promotes anti-proliferative and anti-tumour effects (Paruthiyil *et al.* 2004, Lin *et al.* 2007). It has been proposed that ER β exhibits these effects by decreasing the expression of the E₂-responsive pro-proliferative (c-myc, cyclin A and cyclin D1) (Paruthiyil *et al.* 2004) and anti-apoptotic (Bcl-2

and Bcl-x) (Nilsen *et al.* 2000, Song and Santen 2003) genes, and/or increasing expression of anti-proliferative genes such as p21^{Cip1} and p27^{Kip1} (Paruthiyil *et al.* 2004, Lin *et al.* 2007). In contrast to ER α , the role of ER β in breast cancer cell invasion is not well-studied, but the evidence from at least two studies suggest that ER β inhibits breast cancer cell invasion, both *in vivo* and *in vitro* (Lazennec *et al.* 2001, Thomas *et al.* 2012). Furthermore, studies have shown that ER β can antagonise ER α -mediated effects by either interfering with the recruitment of co-activators required for ER α -mediated actions, or by forming a heterodimer with ER α and thereby affecting the recruitment of this receptor to the promoters of E₂-responsive genes such as the PR (Fox *et al.* 2008).

The presence of ER β in ER α -negative breast cancers is, however, not considered favourable. Studies indicate that in the absence of ER α , ER β promotes proliferation by increasing the expression levels of the proliferative markers Ki67 (O'Neill *et al.* 2004, Skliris *et al.* 2006) and cyclin A (Fox *et al.* 2008). Although the molecular mechanism underlying this phenomenon is unclear, at least one study in the literature suggest that the phosphorylation status of ER β may play a role (Tremblay *et al.* 1999). These authors showed that the phosphorylation of residues within the AF-1 domain in ER β , due to an increase in mitogen-activated protein kinases (MAPK) activity, enhances the recruitment of the co-activator SRC-1, and thereby leads to enhanced transcriptional activity of ER β (Tremblay *et al.* 1999).

1.5. Role of androgens and the AR in breast cancer biology

1.5.1. Androgens

For many decades, E₂ and the ERs were thought to be the main etiological factors implicated in breast cancer pathogenesis. However, recent evidence suggests that androgens and the AR are not only role players in prostate cancer (Weller *et al.* 2014, Karantanos *et al.* 2015), but may also play a role in the development and progression of breast cancer (Nicolás *et al.* 2007, Hu *et*

et al. 2011, Robinson *et al.* 2011, Kotsopoulos and Narod 2012, Li *et al.* 2013). The reported function of androgens in breast cancer is, however, contradictory, as at least one study shows stimulatory effects (Birrell *et al.* 1995), while a number of other studies indicate inhibitory effects on the growth of breast cancer cells (Hackenberg *et al.* 1991, Birrell *et al.* 1998, Greeve *et al.* 2004). Stimulatory effects are supported by studies showing that the most potent natural androgen, 5 α -dihydrotestosterone (DHT), has pro-proliferative effects in MCF-7 and MDA-MB-453 breast cancer cell lines (Birrell *et al.* 1995, Lin *et al.* 2009), suggesting an increased risk of breast cancer. In contrast, the apoptotic and anti-proliferative effects of the potent synthetic androgen, mibelerone (MIB) in T47-D and ZR-75-1 cells, may provide a mechanism to support its reported protective role in breast cancer pathogenesis (Nicolás Díaz-Chico *et al.* 2007, Dimitrakakis 2011, Kotsopoulos and Narod 2012, Coss *et al.* 2014, Lakis *et al.* 2014). Whether androgens elicit anti-proliferative or proliferative effects in cell lines appear to be dependent on various factors including, the specific cell line and androgen used, the concentration of the androgen used, as well as the absence or presence of E₂ (Birrell *et al.* 1995, Birrell *et al.* 1998).

In addition to the above-mentioned *in vitro* studies, both *in vivo* and clinical studies (Dauvois and Labrie 1989, Dauvois *et al.* 1989, Lin *et al.* 2009, Dimitrakakis 2011) have also shown that androgens can either inhibit or stimulate the growth of tumours in breast cancer patients (Sourla *et al.* 1998, Dimitrakakis *et al.* 2003). Clinical evidence suggests that the subcutaneous administration of testosterone in postmenopausal women may be beneficial as it reduces the incidence of breast cancer (Bitzer *et al.* 2008, Dimitrakakis and Bondy 2009, Glaser and Dimitrakakis 2013). The clinical use of testosterone, however, is also associated with negative effects. This is due to the aromatization of testosterone to E₂, thereby possibly increasing the risk of breast cancer (Cauley *et al.* 1999, Somboonporn and Davis 2004). Studies have suggested that DHT, which unlike testosterone cannot be aromatized to E₂, may reduce tumour growth. However, this effect depends on factors such as androgen concentration and cell line

used. Although some studies suggest that DHT may provide a new treatment option for breast cancer patients as it cannot be aromatized to E₂ (Ando *et al.* 2002, Chottanapund *et al.* 2013), DHT may have masculinizing side-effects and thus newly-developed selective androgen receptor modulators (SARMs) may be a better treatment option (Gao and Dalton 2007, Bhasin and Jasuja 2009).

1.5.2. Androgen receptor (AR)

At the molecular level, androgens mediate their effects via the AR (Birrell *et al.* 1998, Kandouz *et al.* 1999, Birrell *et al.* 2007). This intracellular steroid receptor is predominantly cytoplasmic in its ligand-free state and upon androgen binding, the AR translocates to the nucleus where, like the ER, it either positively or negatively regulates target gene expression (Gobinet *et al.* 2002, Lee and Chang 2003). Some AR regulated genes such as the prostate homeobox (NKX3) and kallikreins (KLK5 and KLK3, also known as prostate-specific antigen (PSA)), are associated with breast cancer pathogenesis. For example *in vivo* studies have shown that NKX3 inhibits ER signalling and therefore can be a potential regulator of hormone response in breast cancer (Corthals *et al.* 2000, Xu *et al.* 2001, Nelson *et al.* 2002, Holmes *et al.* 2008, Wang *et al.* 2009, Asch-Kendrick *et al.* 2014). Although PSA is generally regarded as a prostate cancer-specific gene, heightened PSA levels have also been observed in malignant breast cancers. The physiological relevance of the latter is not yet clear, although it has been suggested that PSA promotes breast cancer tumour progression and metastasis (Yu *et al.* 1998, Black *et al.* 2000).

As mentioned earlier, a number of studies have suggested that the AR, which is expressed in approximately 70-90% of primary breast tumours (Søreide *et al.* 1992, Garay and Park 2012) and in 25% of advanced breast tumours (Søreide *et al.* 1992, Birrell *et al.* 1998, Garay and Park 2012, Chottanapund *et al.* 2013), may play a critical role in breast cancer biology (Hu *et al.* 2011, Robinson *et al.* 2011, Li *et al.* 2013). It is proposed that the expression of the AR in breast

tumours inhibits tumour growth and metastasis of cancer, and as a result reduces the number of breast cancer deaths (Søreide *et al.* 1992, Birrell *et al.* 1995, Kulshreshtha *et al.* 2010, Chanplakorn *et al.* 2011, Loibl *et al.* 2011, Yu *et al.* 2011, McNamara *et al.* 2014b). Considering that the majority of breast cancers express the AR, and taking into account that the above-mentioned studies suggest that the AR inhibits breast cancer progression, it has been proposed that the current clinical classification of breast cancers, which focusses only on ER α , PR and human epidermal growth factor receptor 2 (HER2) status, should be altered to include the AR (Farmer *et al.* 2005, McNamara *et al.* 2014b). For example, basal (ER α -negative, AR-negative); luminal (ER α -positive, AR-positive); molecular apocrine (ER α -negative, AR-positive) (Farmer *et al.* 2005). The latter subtype is often referred to as triple negative breast cancer (TNBC) due to the lack of ER α , PR and HER2 expression (Nicholas *et al.* 2003, Anestis *et al.* 2015).

Luminal cancer is subdivided into luminal A (low Ki67 index, slow tumour growth) and B (high Ki67 index, high tumour growth) (Yanagawa *et al.* 2012), with the AR more abundantly expressed in luminal A tumours (Park *et al.* 2010, Loibl *et al.* 2011, Tsang *et al.* 2014). The co-expression of AR with ER is associated with smaller tumour size and no signs of relapse during endocrine treatment and chemotherapy (Castellano *et al.* 2010, Hu *et al.* 2011, Park *et al.* 2011, Tsang *et al.* 2014). The protective effect of the AR is attributed to its ability to inhibit the activity of ER α in patients with luminal breast tumours (Peters *et al.* 2009, Castellano *et al.* 2010). The role of the AR in TNBC is controversial. A study has shown that the expression of the AR in TNBC is associated with an increase in lymph node metastases and number of breast cancer deaths (Hu *et al.* 2011). This is thought to be due to the AR acting as a surrogate ER α and activating transcription of ER α -regulated genes thereby promoting tumour progression in patients (Robinson *et al.* 2011). However, other studies indicate that the presence of the AR in TNBC results in a reduced proportion of lymph node metastases and a better prognosis (Sutton *et al.* 2012, Tang *et al.* 2012). Furthermore, at least one study has shown that AR expression has no influence on TNBC (Mrklič *et al.* 2013).

In conclusion, although studies in the literature on the effects of androgens and the AR on breast cancer are contradictory, the AR is currently being considered as a therapeutic target for the treatment of breast cancer (Chottanapund *et al.* 2013). In ER α -positive (luminal) breast cancers, targetting AR activation may be ideal as the unliganded- or ligand-bound AR inhibits the transcriptional activity of ER α (Peters *et al.* 2009, Fioretti *et al.* 2014). However, in ER α -negative (molecular apocrine/TNBC) breast cancers, it may be beneficial to inhibit the actions of the AR as it may function as a surrogate ER α in the absence of the latter (Fioretti *et al.* 2014). From the above, it is clear that the function of the AR in breast cancer is determined by whether or not the ER is present. In the next section, we will thus discuss the current knowledge in the literature on the association between the ER subtypes and the AR in breast cancer.

1.5.3. Crosstalk between the ER subtypes and the AR

A number of clinical studies have showed that AR expression in ER-positive breast tumours is associated with lower grade disease, reduced invasiveness and longer disease-free survival (Castellano *et al.* 2010, Hu *et al.* 2011, Loibl *et al.* 2011, Park *et al.* 2011, Yu *et al.* 2011, Park *et al.* 2012, Qu *et al.* 2013, Vera-Badillo *et al.* 2013, Witzel *et al.* 2013, Tsang *et al.* 2014). However, these studies do not distinguish between the two ER subtypes expressed in these breast tumours. To the best of our knowledge, the few *in vitro* studies that have been conducted to gain insight into the interplay between the ER subtypes and AR in breast cancer mainly focused on the ER α subtype.

Figure 1.8 shows the proposed mechanisms by which the AR can either promote or inhibit breast cancer pathogenesis. In ER α -positive breast cancer (figure 1.8A), the AR acts as a competitor of ER α and thus suppresses ER α -mediated tumour growth (Birrell *et al.* 1995, Szelei *et al.* 1997, Lanzino *et al.* 2005, Macedo *et al.* 2006, Peters *et al.* 2009, Hickey *et al.* 2012). The above-mentioned is supported by the results from an *in vitro* study by Peters *et al.* (2009)

showing that in ER α -positive breast cancer cells, the AR inhibits the transactivation function of ER α , as well as ER α -mediated cell proliferation (Peters *et al.* 2009). In ER α -negative breast cancer (figure 1.8B), it is proposed that the AR levels increase and that there are changes in cofactor interactions, resulting in the AR mimicking the function of ER α , thereby sustaining tumour growth (Robinson *et al.* 2011, Hickey *et al.* 2012).

Several mechanisms have been identified whereby the AR inhibits the actions of ER α . For example, it has been shown that the AR can bind directly to EREs (Peters *et al.* 2009, Rizza *et al.* 2014), thereby preventing ER α from interacting with the ERE and as a result inhibiting the transactivation function of ER α (Peters *et al.* 2009, Hickey *et al.* 2012, Lamb *et al.* 2014). Considering that both transactivation and transrepression of ER target genes play a role in the regulation of breast cancer cell survival and proliferation, it would be interesting to know whether the transrepression function of ER α would also be inhibited by the AR. Furthermore, ER α and the AR have been shown to interact with similar co-regulatory molecules (Lanzino *et al.* 2005, Fioretti *et al.* 2014), therefore it is possible that when co-expressed these receptors may compete for these molecules. One such example is the AR associated (ARA70) protein, a known AR co-activator that also acts as an ER α co-activator (Lanzino *et al.* 2005). Lastly, ER α and the AR have similar non-genomic activities. This means that the AR as well as ER α interact with similar accessory proteins such as modulator of non-genomic activity of estrogen receptor (MNAR), which is important for cell membrane actions of these receptors. The ability of the AR to interact with these proteins prevents ER α from associating with these proteins thus leading to ER α being unable to regulate signalling pathways at the cell membrane (Fioretti *et al.* 2014). Unlike, the numerous studies examining the interaction between the AR and ER α in breast cancer, not much is known about the interplay between ER β and the AR.




Figure 1.8 Proposed mechanism of the tumour suppressor vs. oncogenic roles of the AR in breast cancer. (A) The AR can act as a tumour suppressor by competing with ER α for binding to EREs, leading to the suppression of ER α -mediated growth. (B) In the absence of ER α , the AR mimics the actions of ER α thereby becoming an oncogene. The red arrow indicates an increase in AR levels. Adapted from (Hickey *et al.* 2012).

1.6. Conclusion

For over a century it has been appreciated that E_2 and the ER play critical roles in breast cancer biology and their role in breast cancer has been extensively studied. $ER\alpha$ has been reported to promote cell growth and survival, while $ER\beta$ inhibits $ER\alpha$ -mediated cell proliferation. However, emerging evidence suggests that other steroid receptors, like the AR, are also involved in the regulation of breast cancer growth (Birrell *et al.* 1998, Brys 2000, Hartman *et al.* 2012, Shah *et al.* 2013, Lakis *et al.* 2014). Interestingly, the evidence in the literature suggests that the precise function of the AR in breast cancer is dependent on whether or not $ER\alpha$ is present, with little known about the link between the AR and $ER\beta$. Figure 1.9 summarises the current knowledge on the interplay between the ER subtypes and the AR in breast cancer. The illustration shows that both the AR and $ER\beta$ can inhibit cell survival and proliferation, directly and indirectly, by suppressing the activity of $ER\alpha$ (Peters *et al.* 2009, Need *et al.* 2012). Also shown is the fact that not much is known about whether the AR can have an effect on the activity of $ER\beta$. Given that both the AR and $ER\beta$ elicits growth-inhibitory effects in the presence of $ER\alpha$, and that the role of the AR and $ER\beta$ changes to promote breast cancer development in the absence of $ER\alpha$, it is important to elucidate the influence of the AR on the function of $ER\beta$. Furthermore, although it is known that the AR modulates the transactivation function of $ER\alpha$, one cannot exclude the fact that ERs elicit their biological activity via both transactivation and transrepression mechanisms. Thus, it is important to also study the effects of the AR on the transrepression function of the ER subtypes. This is particularly important in the light of the fact that inflammation is known to promote the development and progression of breast cancer (Niu *et al.* 2012, Bhatelia *et al.* 2014, Suman *et al.* 2015), and repression of NF κ B-regulated genes via the ER subtypes potentially reduces the risk of inflammation-induced breast cancer (Visser *et al.* 2013).




Figure 1.9. Interplay between the ER subtypes and the AR in breast cancer. ER α promotes cell survival and proliferation, while ER β antagonizes ER α -mediated cell proliferation. The AR has been shown to have the ability to inhibit cell proliferation both directly and indirectly by suppressing the activity of ER α . – and T denote inhibition of activity, + shows activation of activity, the blue arrows demonstrate that the effect is towards ER α while the purple arrow demonstrates that the effect is towards ER β .

1.7. Aims of the study

The evidence in literature suggests that the AR decreases the transactivation function of ER α and that the decrease is more pronounced in the presence of AR ligand (Peters *et al.* 2009). Furthermore, it has been reported that the AR mimics the function of ER α as it can bind to EREs, but that ER α cannot bind to an ARE. Whether the AR has the same effect on the transactivation function of ER β and whether the latter can transactivate via an ARE, is not well studied. Moreover, considering that ERs have both transactivation and transrepression

functions, studies investigating the influence of the AR on the transrepression function of both ER α and ER β are lacking.

Thus, the main aim of the current study was to investigate the influence of the AR on both the transactivation and transrepression functions of the ER subtypes, with the main focus being on ER β . The ER- and AR-deficient MDA-MB-231 human breast cancer cell line was used as an *in vitro* model system for breast cancer to investigate the following:

1. The validation of the experimental system used by Peters *et al.* (2009).
2. The effect of increasing concentrations of human AR, in the absence and presence of AR ligands, on the transactivation function of human ER β via a synthetic ERE-containing promoter reporter construct.
3. The effect of increasing concentrations of the AR on the transactivation function of both ER subtypes on an endogenous ER-regulated gene.
4. The ability of both ER subtypes to transactivate via a synthetic and an endogenous ARE-containing promoter.
5. The effect of the AR on the transrepression function of both ER subtypes via a synthetic NF κ B-containing promoter.
6. The ability of the AR to modulate ER α - or ER β -mediated cell proliferation.

Chapter 2

Materials and

Methods

2.1. Plasmids

The plasmids expressing the human ER α (pSG5-ER α) and ER β (pSG5-ER β) were received from Prof. Frank Gannon (European Molecular Biology Laboratory, Heidelberg, Germany) (Gilles *et al.* 2000). A plasmid expressing the human AR, pSVAR α , (Kastner *et al.* 1990) was obtained from Prof Frank Claessens (University of Leuven, Belgium) (Brinkmann *et al.* 1989). The promoter-reporter construct vit-ERE-TATA-luciferase containing two copies of the estrogen response element (ERE) previously described by (Hall *et al.* 2002) was obtained from Prof Ken Korach (National Institute of Environmental Health Sciences, North Carolina, USA), while the p(IL6 κ B)₃50hu.IL6Pluc⁺ promoter reporter construct was obtained from Prof. Guy Haegeman (University of Gent, Belgium) (Plaisance *et al.* 1997). The pGL2 basic vector, purchased from Promega (Madison, USA), was used as a filler plasmid and the pTAT-GRE-E1b-luc, driven by the E1b promoter containing two copies of the rat TAT-GRE, was obtained from Dr. G. Jenster (Erasmus University of Rotterdam, Netherlands) (Sui *et al.* 1999). The latter promoter-reporter construct will be referred to as an ARE-luc.

2.1.1. Plasmid DNA preparation

Escherichia coli DH5 α cells were made competent using the calcium chloride method and transformed using the heat shock method (Sambrook *et al.* 1989). A starter culture was prepared by growing a single colony of the transformed bacteria in 5 ml Luria-Bertani (LB) medium in the presence of 50 μ g/ml ampicillin (all plasmid vectors used in this study (mentioned in 2.1) contain an ampicillin-resistant gene) for 6 hours while shaking at 37°C. Thereafter 5 ml of the starter cultures were transferred into 250 ml fresh LB and 50 μ g/ml ampicillin. The latter was incubated for 16 hours at 37°C to allow cells to reach exponential growth. Plasmid DNA was purified using the PureYield™ Plasmid Maxiprep System (Promega, Madison, USA) according to the manufacturer's instructions. Briefly, the cultures were centrifuged for 10 minutes at 5000

x g and the supernatants discarded. Each pellet was resuspended in 12 ml Cell Resuspension Solution, before lysis in 12 ml Cell Lysis Solution. Lysis was stopped by adding 12 ml Neutralization Solution, followed by centrifugation of the lysate at 14 000 x g for 30 minutes. Two columns were used to purify the plasmid DNA using a vacuum manifold. The PureYield™ clearing column was set on top of the PureYield™ maxi binding column. The lysate was decanted into the PureYield™ clearing column and maximum vacuum was applied. The PureYield™ clearing column was removed after the lysate had passed through to the PureYield™ maxi binding column. The latter was then washed with 5 ml Endotoxin Removing Buffer, followed by 20 ml Column wash while applying maximum vacuum. The column was subsequently dried for 5 minutes under vacuum, and then inserted into a 50 ml tube. Nuclease-free water was added to the column prior to centrifugation in a swing bucket rotor at 2000 x g for 5 minutes. The eluate containing the plasmid DNA was collected and the DNA concentration determined using a Nanodrop 1000 (Thermo Fisher Scientific, South Africa). Restriction enzyme digests and agarose gel electrophoresis were used to determine the size and integrity (quality) of the plasmid DNA.

2.2. Test compounds

The test compounds used in this study included 17β-estra-1,3,5(10)-etriene-3,17 diol [17-β-estradiol; (E₂)], 5α-androstan-17β-ol-3-one [5α-dihydrotestosterone; (DHT)], 6α-methyl-17α-hydroxyprogesterone acetate [medroxyprogesterone acetate; (MPA)] and 17α-ethynyl-19-nortestosterone 17β-acetate [norethisterone acetate; (NET-A)]. All the test compounds and phorbol 12-myristate 13-acetate (PMA) were obtained from Sigma-Aldrich, South Africa. Test compounds were prepared in absolute ethanol (EtOH) and stored in glass vials at -20°C.

2.3. Cell culture

The human breast cancer cell line, MDA-MB-231 (a kind gift from Prof. Guy Haegemann, University of Gent, Belgium), was maintained in Dulbecco's Modified Eagle's Medium (DMEM) (Sigma-Aldrich, South Africa) containing 4.5 g/ml glucose, phenol-red and supplemented with 10% (v/v) fetal calf serum (FCS) (Separations, South Africa), 100 IU/ml penicillin (Sigma-Aldrich, South Africa) and 100 µg/ml streptomycin (Sigma-Aldrich, South Africa). COS-1 cells (African monkey kidney fibroblast cells), obtained from the American Type Culture Collection (ATCC, Virginia, USA), were maintained in DMEM containing phenol red supplemented with 10% (v/v) FCS, penicillin (100 IU/ ml) and streptomycin (100 µl/ml). All the cell lines were maintained in 75 cm² flasks (Greiner Bio-One International, Austria) at 37°C with a relative humidity of 90% and 5% CO₂. The cell lines were regularly tested for mycoplasma infection using Hoechst staining (Freshney 1987), and only mycoplasma negative cells were used in experiments.

2.4. Promoter reporter assays

2.4.1. Transactivation

MDA-MB-231 cells were seeded into 96-well plates (Porvair Science, United Kingdom) at a density of 1.75×10^4 cells per well in DMEM containing phenol red, 10% (v/v) FCS, 100 IU/ml penicillin and 100 µg/ml streptomycin. The cells were then incubated for 24 hours to allow complete adherence to the surface of the plates. The following day, the cells were rinsed once with sterile 1x PBS and the medium was replaced with phenol red-free DMEM supplemented with 5% (v/v) charcoal treated FCS, 100 IU/ml penicillin and 100 µg/ml streptomycin. Phenol-red free DMEM was used because it is well-known that phenol-red has estrogenic activity (Welshons *et al.* 1988). The cells were subsequently transiently transfected with either 100 ng of the vit-ERE-TATA-luciferase construct and 2.5 ng of the expression vector for human ERα (pSG5-hERα) or 50 ng vit-ERE-TATA-luciferase construct and 2.5 ng of the expression vector

for human ER β (pSG5-hER β), in the absence or presence of 2.5, 5 or 10 ng of the expression vector for human AR (pSVAR_o), using X-tremeGENE HP DNA transfection reagent (Roche Molecular Biochemicals, South Africa) according to the manufacturer's instructions. The filler plasmid pGL2 basic was used to ensure a constant total DNA concentration of 112.5 ng. Transfection and subsequent treatment conditions were similar to that described in Peters *et al.* (2009). Briefly, 3 hours after transfection, the cells were incubated with 0.1% (v/v) EtOH (control) or 1 nM E₂, DHT, MPA or NET-A (agonist mode), or 1 nM E₂ in the absence or presence of 1 nM DHT, MPA or NET-A (antagonist mode) for 36 hours. The cells were subsequently washed with sterile 1x PBS and lysed by adding 25 μ l passive lysis buffer (0.2% (v/v) Triton-X 100, 10% (v/v) glycerol, 2.8% (v/v) Tris-phosphate-ethylenediaminetetra-acetic acid (EDTA) and 1.44 mM EDTA) per well. The plates were then placed on a shaker for 15 minutes at room temperature, before being frozen at -20°C, as the freeze-thawing facilitates effective lysis. On thawing of samples, 10 μ l of the lysates were assayed for luciferase activity using the Luciferase Assay system (Promega luciferase assay reagent, Madison, USA) and a Veritas microplate luminometer (Turner biosystems, USA). In addition, 5 μ l of the cell lysate was used to determine the protein concentration by the Bradford protein determination (Bradford 1976). The luciferase values obtained for all experiments (expressed as relative light units (RLUs)) were normalised relative to the protein concentration for each sample. Some results are expressed as fold induction with the EtOH value set as one, while other results are expressed as the fold induction value in the presence of 1 nM E₂ set as 100% with all other values calculated relative to that. The reported values are averages of at least 3 independent experiments, with each condition performed in triplicate.

2.4.2. Transrepression

The MDA-MB-231 cell line was seeded into 96-well plates at a density of 1.75×10^4 cells per well in DMEM containing phenol-red. After 24 hours, the cells were transiently transfected with 2.5 ng of either pSG5-hER α or pSG5-hER β and 100 ng p(IL6 κ B)₃50hu.IL6Pluc⁺, in the absence or presence of 2.5, 5 or 10 ng pSVAR_o, using the X-tremeGENE HP DNA transfection reagent (Roche Molecular Biochemicals, South Africa) according to the manufacturer's instructions. The filler plasmid, pGL2 basic, was used to ensure a constant total DNA concentration of 112.5 ng. After 3 hours the cells were incubated with 0.1 % (v/v) EtOH in the absence or presence of 10 ng/ml PMA. For agonist mode, cells were treated with 10 ng/ml PMA in the absence (0.1% (v/v) EtOH) or presence of 1 nM E₂, DHT, MPA or NET-A for 36 hours. For antagonist mode, cells were treated with 1 nM E₂ in the absence and presence of 1 nM DHT, MPA or NET-A for 36 hours. Cells were lysed and analysed as in 2.4.1. The luciferase values obtained for all experiments were normalized relative to the protein concentration for each sample. Results are expressed as either the fold induction with the EtOH value set as one in order to quantify the extent of PMA induction, or as normalised percentage response with the value for EtOH + PMA set as 100% in order to determine the extent of repression obtained.

2.5. Western blot analysis

To determine the steroid receptor content in the MDA-MB-231 human breast cancer cell line, cells were plated into 6-well plates (Corning Costar, United Kingdom) at a density of 2.5×10^5 cells per well. For positive controls, COS-1 cells were seeded into 12-well plates (Corning Costar, United Kingdom) at a density of 1×10^5 cells per well and transiently transfected with 250 ng of the respective steroid receptor expression vector for human ER α , ER β or AR (pSG5-ER α , pSG5-ER β or pSVAR_o). Untransfected COS-1 cells were used as negative controls. After 48 hours, the cells were washed with 1x PBS and placed on ice, before lysis with 80 μ l (6 well

plate) or 100 µl (12 well plate) 2x sodium dodecyl sulphate (SDS) sample buffer (1 M Tris-hydrogen chloride (HCl) (pH 6.8), 10% (w/v) SDS, 0.1% (w/v) bromophenol blue, 2% (v/v) β-mercaptoethanol and 20% (v/v) glycerol) (Sambrook *et al.* 1989). Cells were removed from the wells using a cell scraper and then transferred into a 1.5 ml Eppendorf tube (Eppendorf AG, Hamburg, Germany). All lysates were denatured by boiling for 10 minutes at 97°C. Protein samples were separated on a 10% SDS-polyacrylamide gel by performing SDS-polyacrylamide gel electrophoresis (SDS-PAGE) at 100 V for 15 minutes and 200 V for 45 minutes using 1x SDS running buffer (3.5 mM SDS, 25 mM Tris-HCl, 192 mM Glycine). After electrophoresis, the samples on the acrylamide gel were transferred onto a hybond Enhanced Chemiluminescence (ECL) nitrocellulose membrane (Amersham Biosciences, USA) at 180 mA for 90 minutes using 1x transfer buffer (25 mM Tris-HCl (pH 7), 192 mM glycine and 10% methanol). The membranes were blocked for 2 hours in 10% (w/v) fat-free milk powder in 1x TBS (Tris Buffered Saline - 50 mM Tris-HCl and 150 mM NaCl) containing 0.1% (v/v) Tween-20 (TBST). Thereafter, the membranes were rinsed with 1x TBST and incubated with specific primary antibodies directed against the steroid receptors of interest (Santa Cruz Biotechnology Inc, Europe), for 16 hours at 4°C. Glyceraldehyde-3-phosphate dehydrogenase (GAPDH) was used as a loading control. The membranes were then washed with TBST (once for 15 minutes and three 5 minute washes), before incubating with horseradish peroxidase (HRP)-conjugated secondary antibodies (goat anti-rabbit or anti-mouse) (Santa Cruz Biotechnology Inc., Europe) at room temperature for 90 minutes. The dilutions for all antibodies used in this study are summarised in Table 2.1. The membranes were washed again as indicated above, and then visualized using ClarityTM ECL western blotting substrate (Biorad, South Africa) and medical X-ray film (Africa X-Ray Industrial and Medical (Pty) LTD., South Africa).

Table 2.1 Primary and secondary antibody dilutions used for Western blotting

Primary antibody	Dilution	HRP-conjugated secondary antibody	Dilution
ER α (M-20)	1:500	goat anti-rabbit	1:1 000
ER β (H-150)	1:1 000	goat anti-rabbit	1:1 000
AR (441)	1:2 000	goat anti-mouse	1:5 000
GR (H-300)	1:4 000	goat anti-rabbit	1:5 000
MR (H-300)	1:500	goat anti-rabbit	1:5 000
PR-A/B (AB-52)	1:500	goat anti-rabbit	1:2 000
GAPDH (0411)	1:4 000	goat anti-mouse	1:15 000

2.6. Total RNA isolation and complementary DNA (cDNA) synthesis

Human MDA-MB-231 cells were seeded into 12-well plates at a density of 1×10^5 cells per well. Cells were transfected with 33.3 ng of either pSG5-hER α or pSG5-hER β expression vectors in the absence and presence of increasing concentrations of the pSVAR_o (0, 33.3, 66.6 and 133.2 ng, respectively) and pGL2 basic expression vectors to ensure a constant total DNA concentration of 166.5 ng, using X-tremeGENE HP DNA transfection reagent (Roche Molecular Biochemicals, South Africa) according to the manufacturer's instructions. Twenty four hours later the cells were treated for 12 hours with either 0.1% (v/v) EtOH, 1 nM E₂ or DHT, or equimolar amounts of both E₂ and DHT in phenol red-free DMEM supplemented with 5% (v/v) charcoal treated FCS. Following a single PBS wash, the total RNA was isolated from the cells using Tri-reagent (Sigma-Aldrich, South Africa) according to the manufacturer's instructions. Briefly, the cells were lysed by adding 400 μ l Tri-reagent and stored overnight at -80°C, as the freeze-thawing facilitates lysing of the cells. On thawing, the lysates were

transferred to 1.5 ml microcentrifuge tubes and incubated for 5 minutes at room temperature. Thereafter 80 μ l chloroform was added and the samples were vortexed for 1 minute. The samples were incubated for 3 minutes at room temperature before centrifugation at 20 800 x g for 15 minutes at 4°C. The aqueous phase was transferred to a clean microcentrifuge tube and an equal volume of ice-cold isopropanol was added. Samples were vortexed for 1 minute and incubated for 15 minutes at room temperature, followed by centrifugation at 20 800 x g for 15 minutes at 4°C to pellet the RNA. The pellets were subsequently washed by adding 500 μ l 75% (v/v) EtOH, vortexing for 1 minute, and centrifugation at 6800 x g for 5 minutes at 4°C. The supernatant was aspirated and the RNA pellets were air-dried for 10 minutes on ice before dissolving pellets in 15 μ l diethyl pyrocarbonate (DEPC)-treated water. The RNA concentration was measured using a Nanodrop ND-100 Spectrophotometer (Inqaba Biotec, South Africa) and the $A_{260/280}$ ratios were determined. All RNA samples were pure as indicated by $A_{260/280}$ ratios between 1.9 and 2.1 in a 10 mM Tris buffer (Sambrook *et al.* 1989). The integrity of the RNA was confirmed on 1% (w/v) denaturing formaldehyde agarose gels by the presence of intact 28S and 18S ribosomal bands. All RNA samples were stored at -80°C. The total RNA extracted from the transfected MDA-MB-231 cell line was reverse transcribed using the transcriptor first strand cDNA synthesis kit (Roche Applied Science, South Africa) according to the manufacturer's instructions. Briefly, 0.5 μ g total RNA was added to 2.5 μ M anchored-oligo(dT)₁₈-primer and if needed, PCR-grade water was added to a final volume of 6.5 μ l. The template-primer mixture was incubated at 65°C for 10 minutes to denature the RNA secondary structures and immediately placed on ice. Thereafter the following was added: 2 μ l of 1x transcriptor reverse transcriptase reaction buffer (8 mM MgCl₂), 0.25 μ l of protector RNase inhibitor (20 U/ μ l), 1 μ l of deoxynucleotide mix (1 mM per nucleotide) and 0.25 μ l of transcriptor reverse transcriptase (10 U/ μ l) (Roche Applied Science, South Africa). The samples were then incubated at 50°C for an hour in order for reverse transcription to take place.

The reaction was stopped by incubating the samples at 85°C for 5 minutes in order to inactivate the transcript reverse transcriptase. The cDNA samples were stored at –20°C.

2.7. Realtime quantitative polymerase chain reaction (qPCR)

Realtime qPCR was performed using the StepOne Plus Real Time PCR system (Applied Biosystems, South Africa) according to the manufacturer's instructions. The qPCR reaction mixture was prepared by adding a final concentration of 0.2 µM forward and reverse primers respectively (table 2.2), 5 µl 1x SYBR Green (Kapa biosystems, South Africa) and PCR-grade water to a final volume of 9 µl per sample. Thereafter, 1 µl cDNA template or PCR-grade water (negative control) was added. The samples were incubated for 3 minutes at 95°C to activate the Taq polymerase, followed by denaturation for 15 seconds at 95°C. The thermal cycling profiles for PR, PSA (prostate specific antigen) and GAPDH genes are summarized in table 2.3. Agarose gel electrophoresis and melting curve analysis were used to confirm the presence of a single amplicon. A cDNA dilution series was used to generate a standard curve to determine the amplification efficiency of each primer set (Addendum). The efficiencies were 1.96, 1.94 and 2.12 for PR, PSA and GAPDH, respectively. The relative transcript levels of target genes were normalized to the relative transcript levels of GAPDH by using the mathematical model described by (Pfaffl 2001).

Table 2.2. Primer sequences for genes investigated in this study.

Gene	Primers (5' – 3')	Strand	Amplicon size	Reference
			(bp)	
PR	CTTAATCAACTAGGCGAGAG	Forward	122	(Kocanova <i>et al.</i> 2010)
	AAGCTCATCCAAGAATACTG	Reverse		
PSA	AGGCCTTCCTGTACACCAA	Forward	133	(Bohrer <i>et al.</i> 2010)
	GTCTTGGCCTGGTCATTTC	Reverse		
GAPDH	TGAACGGGAAGCTCACTGG	Forward	307	(Ishibashi <i>et al.</i> 2003)
	TCCACCACCCTGTTGCTGTA	Reverse		

Table 2.3. Thermal cycling profiles for all genes investigated.

Gene	Annealing		Elongation		Number of thermal cycles
	Temperature (°C)	Time (seconds)	Temperature (°C)	Time (seconds)	
PR	58	30	72	20	40
PSA	60	10	72	10	45
GAPDH	59	10	72	12	28

2.8. Methyl Thiazolyl Tetrazolium (MTT) cell proliferation assay

The MDA-MB-231 cell line was seeded into 96-well plates at a density of 1×10^4 cells per well. The following day, cells were washed with 200 μ l pre-warmed PBS (37°C) followed by the addition of 200 μ l phenol-red free DMEM, supplemented with 10% (v/w) charcoal-stripped FCS. The cells were subsequently transfected with either 2.5 ng pSG5-hER α , or pSG5-hER β or pGL2 basic using X-tremeGENE HP DNA transfection reagent according to the manufacturer's instructions. After 24 hours, the medium was aspirated and the cells were treated for 48 hours with 0.1% (v/v) (EtOH) or 1 nM E₂ in phenol-red free DMEM supplemented with 10% (v/w) charcoal-stripped FCS only. After 48 hours, 50 μ l of the colorimetric MTT solution (3-(4,5-dimethylthiazolyl-2)-2,5-diphenyltetrazolium bromide) (Sigma-Aldrich, South Africa) prepared by dissolving 5 mg/ml MTT in pre-warmed PBS and filter-sterilising the solution, was added to each well together with 150 μ l phenol red free DMEM containing 10% (v/w) charcoal-stripped FCS. The yellow MTT tetrazolium salt is reduced by metabolically active cells to an insoluble purple formazan precipitate that can be solubilized with dimethyl sulfoxide (DMSO) and quantified spectrophotometrically. After a 4 hour incubation with the MTT solution, the medium was aspirated and 200 μ l DMSO was added to each well to solubilize the crystals. The plate was covered with foil and incubated at room temperature for 5 minutes, while shaking. The absorbance was then read at 540 nm in a Power wave 340 microtiter plate reader (Biotek, Winooski, USA). Results were expressed as fold induction with the value for EtOH set as one.

2.9. Data manipulation and statistical analysis

The GraphPad Prism[®] Software version 5.3 was used for data manipulation, graphical representations and statistical analysis. Two-way ANOVA analysis of variance and Bonferroni's (compares all pairs of columns) post-tests was used for statistical analysis of

grouped data. Unpaired t-tests were used for column graphs with only two sets of data. One-way ANOVA analysis of variance with Bonferroni's (compare all pairs of columns) post-tests as well as Newman-Keuls (compare all pairs of columns) post-tests was used for all column graphs with more than two sets of data. Statistically significant differences are indicated by (*, **or ***) to indicate $p < 0.05$, $p < 0.01$ or $p < 0.001$, respectively, whereas no statistical difference, ($p > 0.05$) is indicated by ns. The letters a, b and c are used to indicate statistical significant differences, with values differing from each other being assigned different letters.

Chapter 3

Results

3.1. Validating the experimental system

A study by Peters *et al.* (2009) showed that the unliganded AR decreases the transcriptional activation of ER α , and that in the presence of liganded AR, this decrease was in a concentration dependent manner. This raised the question of whether the AR could decrease the activity of ER β in a similar manner. To address this question, the model system used by Peters and co-workers was firstly validated, by following similar transfection and treatment conditions in the same cell line (see chapter 2, section 2.4.1). Briefly, the ER and AR negative human MDA-MB-231 breast cancer cell line (absence confirmed by western blot in figure 3.1) was transiently transfected with an ER α expression vector and an estrogen response element (ERE)-driven reporter construct containing two copies of the ERE, in the absence or presence of increasing concentrations of an AR expression vector.




Figure 3.1. The ER subtypes and the AR are not expressed in the MDA-MB-231 breast cancer cell line. Whole cell extracts were prepared from the human MDA-MB-231 (MDA) cell line. Untransfected COS-1 cells were used as the negative control (–ve ctrl), COS-1 cells transiently transfected with pSG5-hER α , pSG5-hER β or pSVAR α expression vectors, respectively were used for the positive controls (+ve ctrl). Protein lysates were analysed by western blotting using antibodies specific to ER α , ER β , AR or GAPDH, respectively. GAPDH was used as the loading control.

The cells were subsequently treated with 1 nM E₂, the most biologically active estrogen (Samavat and Kurzer 2015). As expected and shown in figure 3.2A, 1 nM E₂ caused significant

transactivation ($p < 0.001$) of an ERE-containing promoter-reporter construct via ER α (3.8-fold). Consistent with the results shown by Peters and co-workers, the result in figure 3.2B shows that the AR inhibits E₂-induced ER α transactivation. However, it should be noted that the current study shows a concentration-dependent decrease in the presence of the unliganded AR, and that the decrease is more pronounced than the decrease observed by Peters *et al.* (2009). Furthermore, to exclude a mechanism whereby the AR decreases ER α protein expression, we, like Peters and co-workers, used western blot analysis to show that increasing concentrations of the AR do not affect ER α protein levels (figure 3.3).




Figure 3.2 legend on next page.

Figure 3.2. The AR decreases E₂-induced ER α transactivation function. Human MDA-MB-231 breast cancer cells were transiently transfected with pSG5-hER α and the vit-ERE-TATA-luciferase reporter construct, in the absence or presence of increasing concentrations of pSVAR_O (1x, 2x or 4x molar excess), and where needed pGL2basic, using X-tremeGENE HP DNA transfection reagent according to the manufacturer's instructions. After 3 hours cells were treated with 0.1% (v/v) EtOH (control) or 1 nM E₂ for 36 hours. The luciferase values were normalised to the protein concentrations, and expressed as luciferase/protein (luc/prot). The fold induction at 1 nM E₂ via ER α is shown in (A), while in (B) the fold induction in (A) was set as 100%, with induction in the presence of various concentrations of AR calculated relative to that. The results indicate the average (\pm SEM) of at least three independent experiments, with each condition performed in triplicate. Statistical analysis was performed using the unpaired t-test (A) and one-way ANOVA analysis of variance with Bonferroni (compare all pairs of columns) (B) post-tests.




Figure 3.3 legend on next page.

Figure 3.3. The overexpressed AR does not affect the protein levels of overexpressed ER α in the MDA-MB-231 breast cancer cell line. (A) Human MDA-MB-231 breast cancer cells were transiently transfected as in figure 3.2. Protein lysates were analysed by western blotting with antibodies specific to ER α , AR or GAPDH (loading control). Protein lysates from COS-1 cells transfected with pSG5-hER α and pSVAR₀ were used as positive controls (+ve), while lysates from untransfected COS-1 cells were used for the negative control (–ve ctrl). (B) Western blots were quantified using UN-SCAN-IT software.

3.2. The AR does not affect the transcriptional activation of ER β on a synthetic ERE-containing promoter

As the experimental system had been validated, we next investigated the effect of increasing concentrations of the AR on ER β protein levels as well as on ER β transcriptional activation. The human MDA-MB-231 cell line was transiently transfected with 2.5 ng pSG5-ER β and 50 ng vit-ERE-TATA-luciferase in the absence or presence of increasing concentrations of pSVAR₀ (2.5 ng, 5 ng or 10 ng) and where needed, pGL2basic, to ensure that the total amount of DNA transfected remained constant. The cells were subsequently treated with 1 nM E₂ for 36 hours. The western blot in figure 3.4 indicates that, similar to the observation for ER α (figure 3.3), the AR does not affect ER β protein levels. Having shown that the AR does not affect ER β protein levels, the effects of increasing concentrations of the AR on the transcriptional activation of ER β was next investigated. The result in figure 3.5A shows that 1 nM E₂ caused significant transactivation ($p < 0.001$) of an ERE-containing promoter-reporter construct via ER β . However, in contrast to the results obtained for ER α (figure 3.2B), the AR does not affect E₂-induced transcriptional activation via ER β (figure 3.5B).




Figure 3.4. The overexpressed AR does not affect protein levels of overexpressed ERβ in the MDA-MB-231 human breast cancer cell line. (A) Human MDA-MB-231 breast cancer cells were transiently transfected with pSG5-hERβ and the vit-ERE-TATA-luciferase reporter construct, in the absence or presence of increasing concentrations of pSVAR₀ (1x, 2x or 4x molar excess), and where needed pGL2basic, using X-tremeGENE HP DNA transfection reagent according to the manufacturer's instructions. After 3 hours cells were treated with 0.1% (v/v) EtOH (control) or 1 nM E₂ for 36 hours. Protein lysates were analysed by western blotting with antibodies specific to ERβ, AR or GAPDH (loading control). Protein lysates from COS-1 cells transfected with pSG5-hERβ and pSVAR₀ were used as positive controls (+ve), while lysates from untransfected COS-1 cells were used for the negative control (-ve ctrl). (B) Western blots were quantified using UN-SCAN-IT software.




Figure 3.5. The AR does not decrease the transactivation function of E₂-bound ER β . Human MDA-MB-231 breast cancer cells were transfected as in figure 3.4. The luciferase values were normalised to the protein concentrations, and expressed as luciferase/protein (luc/prot). The fold induction at 1 nM E₂ via ER β is shown in (A), while in (B) the fold induction in (A) was set as 100%, with induction in the presence of various concentrations of AR calculated relative to that. The results indicate the average (\pm SEM) of at least three independent experiments, with each condition performed in triplicate. Statistical analysis was performed using the unpaired t-test (A) and one-way ANOVA analysis of variance with Bonferroni (compare all pairs of columns) (B) post-tests.

3.3. The AR ligands do not augment the effect of the unliganded AR

Although we established that the unliganded AR does not affect E₂-bound ER β transcriptional activation, we next investigated whether the ligand-activated AR would modulate ER β activity. This was based on the fact that Peters and co-workers showed that when the AR is activated by its natural ligand, DHT or the selective AR modulator, MPA, a concentration dependent decrease in ER α activity was observed as the AR levels were increased (Peters *et al.* 2009). We also used DHT and MPA for model validation, but included NET-A, a progestin previously reported to have similar potent androgenic activity as DHT and MPA (Africander *et al.* 2014).

The human MDA-MB-231 cells were transiently transfected with 2.5 ng pSG5-ER α and 100 ng vit-ERE-TATA-luciferase or 2.5 ng pSG5-ER β and 50 ng vit-ERE-TATA-luciferase, in the absence or presence of increasing concentrations of pSVARo (2.5 ng, 5 ng, 10 ng). The cells were subsequently treated with 1 nM E₂ in the absence or presence of 1 nM DHT, MPA or NET-A for 36 hours. The results in figure 3.6A-C show that inhibition of ER α transactivation function by the AR is not augmented by the addition of AR ligand. Surprisingly, cells transfected with only ER α and treated with equimolar concentrations of E₂ and MPA or E₂ and NET-A, displayed a significant decrease of E₂ activity via ER α (figure 3.6B-C), suggesting that MPA and NET-A are ER α antagonists. None of the AR ligands displayed agonist activity via ER α (figure 3.6D).




Figure 3.6 continues on next page.




Figure 3.6 legend on next page.

Figure 3.6. Inhibition of ER α transcriptional activation is not enhanced in the presence of AR ligands. Human MDA-MB-231 breast cancer cells were transiently transfected with pSG5-hER α and the vit-ERE-TATA-luciferase reporter construct, in the absence or presence of increasing concentrations of pSVAR_O (1x, 2x or 4x molar excess), and where needed pGL2basic, using X-tremeGENE HP DNA transfection reagent according to the manufacturer's instructions. After 3 hours, cells were treated with 0.1% (v/v) EtOH (control) or 1 nM E₂ in the absence or presence of 1 nM DHT, MPA or NET-A for 36 hours. The luciferase values were normalised to protein concentrations, and expressed as luciferase/protein (luc/prot). The results indicate the average (\pm SEM) of at least three independent experiments, with each condition performed in triplicate. **(A -C)** The fold induction at 1 nM E₂ via ER α was set as 100%, and induction in the presence of various concentrations of liganded AR calculated relative to that. **(D)** The fold induction at 1 nM E₂, DHT, MPA and NET-A via ER α is shown. Statistical analysis was performed using Newman-Keuls (compare all pairs of columns) post-tests (A-C) and one-way ANOVA analysis of variance with Bonferroni (compare all pairs of columns) post-tests (D).

Similar to the results obtained for ER α , results in figure 3.7A-C indicate that the AR ligands do not modulate the effect of unliganded AR on the transactivation function of ER β , and none of the AR ligands displayed agonist activity via ER β (figure 3.7D). Interestingly, in contrast to the results seen for ER α (figure 3.6B-C), neither MPA nor NET-A had an inhibitory effect on the E₂-induced activity of ER β (figure 3.7B-C). Taken together, these results indicate that the inhibition of ER α transactivation function by the AR is independent of AR ligand. Furthermore, the transactivation function of ER β is not influenced by either the unliganded or liganded AR.




Figure 3.7 continues on next page.




Figure 3.7 legend on next page.

Figure 3.7. Ligand bound AR does not decrease the transcriptional activity of E₂ via ER β . Human MDA-MB-231 breast cancer cells were transiently transfected with pSG5-hER β and the vit-ERE-TATA-luciferase reporter construct, in the absence or presence of increasing concentrations of pSVAR_O (1x, 2x or 4x molar excess), and where needed pGL2basic, using X-tremeGENE HP DNA transfection reagent according to the manufacturer's instructions. After 3 hours, cells were treated with 0.1% (v/v) EtOH (control) or 1 nM E₂ in the absence or presence of 1 nM DHT, MPA or NET-A for 36 hours. The luciferase values were normalised to protein concentrations, and expressed as luciferase/protein (luc/prot). The results indicate the average (\pm SEM) of at least three independent experiments, with each condition performed in triplicate. **(A -C)** The fold induction at 1 nM E₂ via ER β was set as 100%, and induction in the presence of various concentrations of liganded AR calculated relative to that, and **(D)** The fold induction at 1 nM E₂, DHT, MPA and NET-A via ER β . Statistical analysis was performed using two-way ANOVA analysis of variance with Bonferroni (compare all pairs of columns) post-tests (A-C) and one-way ANOVA analysis of variance with Bonferroni (compare all pairs of columns) post-tests (D).

3.4. AR abolishes the transcriptional activity of E₂ via both ER subtypes on an endogenous ER regulated gene in a breast cancer cell line

To investigate whether the results observed on the synthetic ERE-containing promoter could be mimicked on an endogenous ER regulated gene, the MDA-MB-21 breast cancer cell line was transiently transfected with 2.5 ng of either the ER α or the ER β expression vector. The cells were treated with 1 nM E₂ for 12 hours, followed by qPCR analysis for the mRNA expression of the PR, a well-known ER regulated gene, which is used as a marker for functional ER signalling (Han *et al.* 2008, Sivik and Jansson 2012). The optimal time for PR gene expression in the presence of both ER subtypes was 12 hours (Addendum). The results in figure 3.8A show that E₂ upregulates PR mRNA expression in MDA-MB-231 cells transfected with either ER α (2.4-fold) or ER β (1.9-fold). As a control to determine whether the E₂ response was indeed via the particular overexpressed ER subtypes, the cell line was transfected with 2.5 ng of the empty vector pGL2basic. The result in figure 3.8A shows that E₂ has no effect on PR gene expression

in the absence of the transfected ER subtypes, suggesting that the E₂-mediated upregulation of the PR gene is ER dependent.

Next we investigated how the presence of increasing AR concentrations would influence the E₂-induced upregulation of the PR gene, by transfecting the MDA-MB-21 breast cancer cell line with 2.5 ng of either the ER α or the ER β expression vector in the absence or presence of 2.5, 5 or 10 ng of an AR expression vector. Note that, as we had shown that the addition of AR ligands in the presence of the AR did not influence ER transactivation on a synthetic ERE-containing promoter, we did not investigate PR gene expression in the presence of the liganded AR. The result in figure 3.8B shows that in the presence of transfected ER α , the unliganded AR appears to decrease the E₂-induced upregulation of the PR gene in a concentration dependent manner. However, the trend indicating that the decrease is concentration-dependent, was not statistically significant. Surprisingly, and in contrast to results on the synthetic ERE-containing promoter, the unliganded AR also appeared to decrease the E₂-induced upregulation of the PR gene in the presence of transfected ER β (figure 3.8C).




Figure 3.8 continues on next page.




Figure 3.8. The E₂-induced expression of the PR gene in MDA-MB-231 breast cancer cells transfected with either ER α or ER β is inhibited by the AR. Human MDA-MB-231 breast cancer cells were transiently transfected with pSG5-hER β or pSG5-hER α in the absence or presence of increasing concentrations of pSVAR_O (1x, 2x or 4x molar excess), and where needed the filler plasmid pGL2basic using X-tremeGENE HP DNA transfection reagent according to the manufacturer's instructions. After 24 hours, cells were treated with 0.1% (v/v) EtOH (control) or 1 nM E₂ for 12 hours. Total RNA was isolated, cDNA synthesized and mRNA expression levels of PR and GAPDH were determined by QPCR. GAPDH was used as the internal standard and the ratio of PR mRNA expression/GAPDH mRNA of samples treated with E₂ were calculated relative to EtOH. The fold induction is shown in (A), while this fold induction at 1 nM E₂ via ER α (B) or ER β (C) was set as 100%, and mRNA expression in the presence of various concentrations of AR calculated relative to that. Results indicate the average (\pm SEM) of four independent experiments. Statistical analysis was performed using one-way ANOVA analysis of variance and Bonferroni (compare all pairs of columns) post-tests.

3.5. Androgen response element transactivation via ER α but not ER β

It is well-known that the ER and AR activate transcription of target genes via binding to distinct response elements namely the ERE for ER (Kumar and Chambon 1988, Hyder *et al.* 1999, Ishida *et al.* 2010, Botelho *et al.* 2015), and the androgen response element (ARE) for AR (Mangelsdorf *et al.* 1995, Mitchell *et al.* 2000, Haendler *et al.* 2001, Matsumoto *et al.* 2013). However, recent evidence suggests that while ER α only binds to its consensus ERE, the AR can also bind to an ERE (Peters *et al.* 2009). Using promoter reporter assays, we thus next investigated whether either ER α or ER β could transactivate via an ARE. MDA-MB-231 cells were transfected with 2.5 ng of either pSG5-ER α or pSG5-ER β and 100 ng of the pTAT-GRE-E1b-luc promoter-reporter construct. As most steroid receptors, except the ER (Klock *et al.* 1987, Klein-Hitpass *et al.* 1988, Brosens *et al.* 2004), can bind the same DNA response element (Beato *et al.* 1989), the GRE can also serve as a response element for the AR, and is then referred to as an ARE. Results in figure 3.9A indicate that after treatment with 1 nM E₂ or DHT ER β does not transactivate via an ARE ($p > 0.05$). Considering that Peters *et al.* (2009) showed that ER α does not bind to an ARE, we were surprised when our results indicated significant transactivation ($p < 0.001$) of an ARE-containing promoter-reporter construct via ER α (figure 3.9B). In addition, significant transactivation was also observed in the presence of 1 nM DHT (figure 3.9B).

As a control to determine whether the ARE-containing promoter-reporter construct is transcriptionally active, cells were also transfected with the pSVARo expression vector and the ARE-containing promoter-reporter construct, and treated with 1 nM DHT or E₂. As expected, induction (317-fold) was observed when cells transfected with the AR and an ARE were treated with 1 nM DHT (figure 3.9C), indicating that DHT and the ARE promoter-reporter construct are active in this cell line. No significant induction was observed with 1 nM E₂ indicating that E₂ has no agonist activity via the AR on an ARE.




Figure 3.9 legend on next page.

Figure 3.9. Unlike ER α , ER β does not transactivate via a synthetic ARE promoter-reporter construct. Human MDA-MB-231 breast cancer cells were transiently transfected with (A) pSG5-hER β , (B) pSG5-hER α or (C) pSVAR₀ and a pTAT-GRE-E1b-luciferase containing promoter-reporter construct using X-tremeGENE HP DNA transfection reagent according to the manufacturer's instructions. After 3 hours, cells were treated for 36 hours with 0.1% (v/v) EtOH (control) or 1 nM E₂ or DHT. Luciferase values were normalised to the protein concentrations, and expressed as luciferase/protein (luc/prot). The fold induction via ER β (A), ER α (B) and the AR (C) on an ARE is shown. The results indicate the average (\pm SEM) of two independent experiments, with each condition performed in triplicate. Statistical analysis was performed using one-way ANOVA analysis of variance and Bonferroni (compare all pairs of columns) post-tests.

3.6. E₂-induced upregulation of an endogenous ARE-containing gene can be mediated by both ER α and β

To investigate whether the results observed on the synthetic ARE-containing promoter could be mimicked on an endogenous ARE-containing promoter, the MDA-MB-21 breast cancer cell line was transiently transfected with 2.5 ng of either the ER α or the ER β expression vector. The cells were treated with 1 nM E₂ or DHT for 12 hours, followed by qPCR analysis for the mRNA expression of the prostate specific antigen (PSA), which is a well-known ARE-containing gene. In contrast to the results in figure 3.9A and B, where we observed that unlike ER α , ER β does not transactivate via a synthetic ARE, the results in figure 3.10A and B show that PSA mRNA expression was significantly upregulated in the presence of E₂ in cells transfected with either ER β (6.0-fold) or ER α (2.6-fold). As expected, results from cells transfected with the AR and treated with 1 nM DHT showed significant upregulation (2.1-fold) of PSA gene expression (figure 3.10C).




Figure 3.10 legend on next page.

Figure 3.10. The E₂-induced mRNA expression of an AR regulated gene (PSA) via both ER α and ER β . Human MDA-MB-231 breast cancer cells were transiently transfected with (A) pSG5-hER β (B) pSG5-hER α or (C) pSVAR₀ using X-tremeGENE HP DNA transfection reagent according to the manufacturer's instructions and incubated for 24 hours. Cells were subsequently treated with 0.1% (v/v) EtOH (control) or 1 nM E₂ or DHT for 12 hours. Total RNA was isolated, cDNA synthesized and mRNA expression levels of PSA and GAPDH were determined by QPCR. GAPDH was used as the internal standard and the ratio of PSA mRNA expression/GAPDH mRNA of all samples were calculated relative to the sample treated with 0.1% (v/v) EtOH. Results show the PSA mRNA expression at 1 nM E₂ or DHT via (A) ER β , or (B) ER α or (C) AR. For A and B results indicate the average (\pm SEM) of two independent experiments, while C is the result of a single experiment. Statistical analysis was performed using one-way ANOVA analysis of variance and Bonferroni (compare all pairs of columns) post-tests (A and B) and the unpaired t-test (C).

3.7. The AR inhibits the transrepression function of both ER α and ER β

As the ER elicits its biological effects by either activating or repressing the transcription of target genes (Harrington *et al.* 2003), this study also examined the influence of the AR on the transrepression function of both ER α and ER β . MDA-MB-231 cells were transiently transfected with 2.5 ng of either pSG5-ER α or pSG5-ER β and 100 ng of a synthetic IL-6NF κ B3-luciferase promoter-reporter construct, in the absence or presence of increasing concentrations of pSVAR₀. The cells were subsequently treated with 10 ng/ml PMA in the absence and presence of 1 nM E₂. As shown in figure 3.11A and C, stimulating the cells with PMA significantly induced an NF κ B driven induction of 3.4-fold and 8.4-fold in the presence of ER α and ER β , respectively. The PMA-induced response was then set to 100% and the E₂ response via ER α (figure 3.11B) or ER β (figure 3.11D) calculated as a percentage relative to that. The E₂-mediated repression on the NF κ B promoter, via ER α and ER β , was 87.8% and 48.5%, respectively. While the addition of increasing concentrations of the AR inhibited the E₂-mediated repression on the NF κ B promoter via ER α (figure 3.11B), E₂-mediated repression via ER β was not affected by the AR (figure 3.11D).




Figure 3.11 continues on next page.




Figure 3.11 legend on next page.

Figure 3.11. The AR inhibits the transrepression function of both ER α and ER β . Human MDA-MB-231 breast cancer cells were transiently transfected with either (A) pSG5-hER α or (B) pSG5-hER β and the IL-6NF κ B3-luciferase containing promoter-reporter construct, in the absence or presence of increasing concentrations of pSVAR₀, (1x, 2x or 4x molar excess), and where needed pGL2basic, using X-tremeGENE HP DNA transfection reagent according to the manufacturer's instructions. After 3 hours, cells were treated with 0.1% (v/v) EtOH (control) in the absence or presence of 10 ng/ml PMA (A and C) or 10 ng/ml PMA in the absence and presence of 1 nM E₂ (B and D) for 36 hours. The luciferase values were normalised to protein concentrations, and expressed as luciferase/protein (luc/prot). The results indicate the average (\pm SEM) of two independent experiments, with each condition performed in triplicate. (A and C) EtOH+PMA induction via the NF- κ B-containing promoter construct calculated relative to the EtOH set as 1. (B and D) The PMA-induced response was set as 100% and the E₂-induced transrepression via ER α or ER β , in the absence or presence of AR, was calculated relative to that. Statistical analysis was performed using unpaired t-test (A and C) and one-way ANOVA analysis of variance with Bonferroni (compare all pairs of columns) post-tests (B and D).

It was next investigated whether the presence of an AR ligand could modulate these effects. The human MDA-MB-231 cells were transfected as above and were subsequently treated with 1 nM E₂ in the absence or presence of 1 nM DHT, MPA or NET-A for 36 hours. Similar to results in figure 3.6 showing that the inhibition of ER α transactivation function by the AR is not enhanced by the presence of AR ligands, results in figure 3.12A-C indicate that AR ligands do not influence the AR-dependent inhibition of ER α transrepression function. Surprisingly, although the unliganded AR had no effect on the transrepression function of ER β , results in figure 3.13A-C indicate that the ligand-bound AR inhibits the transrepression function of ER β . This result is in contrast to the results in figure 3.7 showing that both the unliganded and liganded AR had no effect on the transactivation function of ER β .




Figure 3.12 continues on next page.





Figure 3.12. Inhibition of ER α transcriptional repression is not enhanced by the presence of AR ligands. Human MDA-MB-231 breast cancer cells were transiently transfected with pSG5-hER α and the IL-6NF κ B3-luciferase containing promoter-reporter construct, in the absence or presence of increasing concentrations of pSVAR_O (1x, 2x or 4x molar excess), and where needed pGL2basic, using X-tremeGENE HP DNA transfection reagent according to the manufacturer's instructions. After 3 hours, cells were treated with 10 ng/ml PMA and 1 nM E₂ in the absence or presence of 1 nM DHT, MPA or NET-A for 36 hours. The luciferase values were normalised to protein concentrations, and expressed as luciferase/protein (luc/prot). The results indicate the average (\pm SEM) of two independent experiments, with each condition performed in triplicate. The E₂-induced transrepression on the NF κ B-containing promoter construct via ER α , in the absence or presence of AR, was expressed as a percentage calculated relative to EtOH + PMA set as a 100%. Statistical analysis was performed using one-way ANOVA analysis of variance with Bonferroni (compare all pairs of columns) post-tests as well as two-way ANOVA analysis of variance with Bonferroni (compare all pairs of columns) post-tests.

A



B




Figure 3.13 continues on next page.




Figure 3.13. The ligand-bound AR inhibits the E₂-induced transrepression function of ERβ. Human MDA-MB-231 breast cancer cells were transiently transfected with pSG5-hERβ and the IL-6NFκB3-luciferase containing promoter-reporter construct, in the absence or presence of increasing concentrations of pSVAR_O (1x, 2x or 4x molar excess), and where needed pGL2basic, using X-tremeGENE HP DNA transfection reagent according to the manufacturer's instructions. After 3 hours, cells were treated with 10 ng/ml PMA and 1 nM E₂ in the absence or presence of 1 nM DHT, MPA or NET-A for 36 hours. The luciferase values were normalised to protein concentrations, and expressed as luciferase/protein (luc/prot). The results indicate the average (± SEM) of two independent experiments, with each condition performed in triplicate. The E₂-induced transrepression on the NFκB-containing promoter construct via ERβ, in the absence or presence of AR, was expressed as a percentage calculated relative to EtOH + PMA set as a 100%. Statistical analysis was performed using one-way ANOVA analysis of variance with Bonferroni (compare all pairs of columns) post-tests as well as two-way ANOVA analysis of variance with Bonferroni (compare all pairs of columns) post-tests.

3.8. No change in proliferation was observed in cells transfected with ER

Cell proliferation assays were used to investigate E₂-induced proliferation of the MDA-MB-231 breast cancer cell line when transiently transfected with either ER α or ER β , and how proliferation could be modulated by the presence of overexpressed AR. The MDA-MB-231 cell line was thus transfected with 2.5 ng of the expression vectors for either pSG5-ER α , pSG5-ER β or the pGL2basic empty vector. The latter was used as a control to determine if E₂-induced cell proliferation was indeed via the overexpressed ER subtypes. As expected, no cell proliferation was observed for cells transfected with the empty vector (figure 3.14). Surprisingly, however, no E₂-induced cell proliferation was detected when cells were transfected with either ER α or ER β (figure 3.14), and thus we could not investigate the effects of the AR on ER-mediated cell proliferation.




Figure 3.14 legend on next page.

Figure 3.14. Human MDA-MB-231 breast cancer cells were transiently transfected with pGL2basic (empty vector) or pSG5-hER α or pSG5-hER β or using X-tremeGENE HP DNA transfection reagent according to the manufacturer's instructions. After 24 hours these cells, as well as untransfected MDA-MB-231 cells, were treated for 48 hours with 0.1% (v/v) EtOH (control) or 1 nM E₂. The results indicate the relative fold cell proliferation under various conditions with the value for cells treated with EtOH for each condition (control) set as 1. The average (\pm SEM) of at least four independent experiments, with each condition performed in triplicate is shown. Statistical analysis was performed using one-way ANOVA analysis of variance with Bonferroni (compare all pairs of columns) post-tests.

Chapter 4

Discussion

4.1. Introduction

Steroid receptors play an important role in the pathogenesis of breast cancer and understanding their clinical significance is a crucial aspect of breast cancer research (Welboren *et al.* 2007, Han 2014, Sabatier *et al.* 2014). The ER is the most common and well established breast cancer biomarker, with 75% of breast cancers being ER-positive. The ER exists as two subtypes, ER α and ER β , which are transcription factors that modulate the expression of target genes (Couse and Korach 1999). The activated ER α stimulates breast cancer cell proliferation by regulating specific genes involved in the cell cycle (Frasor *et al.* 2003, Hartman *et al.* 2009), while ER β has been shown to regulate a variety of target genes resulting in ER β opposing the actions of ER α (Hartman *et al.* 2009). For example, E₂-induced ER α stimulates breast cancer cell proliferation by activating cyclin D1, an oncogene which regulates G1-S cell cycle progression, while ER β is anti-proliferative as it inhibits cyclin D1 activation (Paech *et al.* 1997, Weatherman *et al.* 2001, Liu *et al.* 2002). Due to the distinct roles of ER α and ER β in breast cancer, several studies have suggested that blocking the activity of ER α , but not ER β , is an effective treatment option for breast cancer (Assikis and Craig 1997).

Although the role of the ER subtypes in breast cancer pathogenesis has been well-characterized, recent studies are also exploring the role of other steroid receptors (Peters *et al.* 2009, Need *et al.* 2012, Obr and Edwards 2012, Skor *et al.* 2013, Cochrane *et al.* 2014, Knutson and Lange 2014, Pistelli *et al.* 2014, Rechoum *et al.* 2014, Abduljabbar *et al.* 2015). For example, as the AR is expressed in 70% of breast cancer tumours (Birrell *et al.* 1998, Schippinger *et al.* 2006, Lin *et al.* 2009, Hu *et al.* 2011, Chottanapund *et al.* 2013), many clinical (Hu *et al.* 2011, Park *et al.* 2011, Pistelli *et al.* 2014) and molecular studies (Peters *et al.* 2009, Chottanapund *et al.* 2013, Cochrane *et al.* 2014, Rechoum *et al.* 2014) are investigating the role of the AR and androgens in breast cancer. Of particular interest is the fact that AR signalling is known to exert anti-estrogenic effects (Dauvois *et al.* 1991, Ando *et al.* 2002) suggesting crosstalk between the

ER and AR signalling pathways. Indeed, the study by Peters and co-workers showed that both the unliganded and liganded AR has the ability to inhibit the activity of ER α (Peters *et al.* 2009). In agreement with this, two other studies have also reported interplay between ER α and AR transcriptional pathways (Need *et al.* 2012, Rechoum *et al.* 2014). Whether crosstalk also exists between the AR and ER β in breast cancer cells is not well-studied and thus formed the basis of this thesis. In chapter 3, it was thus investigated whether the unliganded and/or ligand-bound AR, could modulate the transcriptional activity (transactivation and transrepression) of ER β in the human MDA-MB-231 breast cancer epithelial cell line.

4.2. Validating the model system

In the first part of this thesis, using western blot analysis (figure 3.1), previous reports in the literature indicating that the human MDA-MB-231 breast cancer cell line does not express the ER subtypes and the AR was confirmed (Horwitz *et al.* 1978, Lazennec and Katzenellenbogen 1999, Leo *et al.* 2004, Buchanan *et al.* 2005, Abduljabbar *et al.* 2015). This result, together with previous reports that the MDA-MB-231 cells do not express the PR, but express the GR and the MR, suggests that the human MDA-MB-231 cell line is an appropriate model cell line for comparing the activities of specific transiently transfected steroid hormone receptors.

Next, the experimental conditions used in the study by Peters *et al.* (2009) were mimicked, and in agreement with their results, the current study showed that the unliganded AR (figure 3.2B) inhibits the transcriptional activation of ER α (figure 3.2B). However, Peters and co-workers showed that the decrease in ER α transcriptional activation increased as the amount of liganded AR, activated by either the natural androgen DHT or the selective AR modulator MPA (MacLaughlin and Richardson 1979, Teulings *et al.* 1980, Birrell *et al.* 1998), was increased. The current study, on the other hand, showed that a concentration-dependent decrease in the transcriptional activation of ER α was observed in the presence of the unliganded AR, and that

the AR ligand (DHT, MPA or NET-A (a progestin with androgenic activity (Africander *et al.* 2014)), did not increase this effect (figures 3.2B; figures 3.6A-C). Furthermore, consistent with the result of Peters *et al.* (2009), the western blot analysis in figure 3.3 clearly indicates that the inhibition of ER α transcriptional activation by the AR was not due to the AR decreasing ER α protein levels.

Furthermore, our results show that MPA and NET-A, but not DHT antagonize the E₂-induced agonist activity for transactivation via ER α (figure 3.6B and C). It is known that progestins that exert androgenic properties, like MPA, are anti-estrogenic (Clarke and Sutherland 1990, Dauvois *et al.* 1991, Ando *et al.* 2002) via a mechanism that does not involve ER binding (Leung *et al.* 1981, Vignon *et al.* 1983, Sutherland *et al.* 1988). For example, it has been shown that MPA decreases intracellular E₂ levels as a result of it enhancing 17 β -HSD2 activity, which causes the conversion of E₂ to E₁ (Dauvois *et al.* 1991, Pasqualini 2003). Considering that NET-A, like MPA, has potent androgenic properties (Africander *et al.* 2014), and that 17 β -HSD type 1 and type 2 are expressed in MDA-MB-231 cells (Day *et al.* 2006, Newman *et al.* 2006), it is likely that the anti-estrogenic activity seen by both MPA and NET-A is due to these ligands increasing 17 β -HSD2 expression and activity.

4.3. Effects on synthetic promoter-reporter genes are not always reflected on endogenous genes

Having validated the model system used by Peters *et al.* (2009) for ER α , we next used a similar model system to investigate the effect of the AR on the transactivation function of ER β . Results in figure 3.5B showed that when MDA-MB-231 cells were co-transfected with the ER β cDNA expression vector and increasing concentrations of the AR expression vector, the unliganded AR did not affect the E₂-induced transcriptional activation of ER β . These findings are consistent with a previous study in CV-1 monkey kidney cells showing that co-expression of

the liganded AR with ER β did not cause a significant decrease in ER β transactivation (Panet-Raymond *et al.* 2000). Furthermore, our investigations in the presence of AR ligands (DHT, MPA, NET-A) showed that, like the unliganded AR, ligand-bound AR has no effect on the transactivation function of ER β on a simple ERE (figure 3.7). Interestingly, using a yeast two-hybrid assay, Panet-Raymond and co-workers showed no interaction between the AR and ER β , but like Peters *et al.* (2009) showed an interaction between the AR and ER α (Panet-Raymond *et al.* 2000, Peters *et al.* 2009). Interestingly, these observations occurred in the presence of either unliganded or liganded receptors. Several molecular mechanisms have been proposed on how the AR suppresses the transactivation function of ER α (Fioretti *et al.* 2014). For example, the AR and ER α have been shown to directly interact with each other thus preventing the latter from binding to EREs and as it has also been shown that the AR can bind to both synthetic and endogenous EREs (Peters *et al.* 2009, Fioretti *et al.* 2014), it can compete with ER α for binding to the EREs.

We next determined the effects of increasing amounts of unliganded AR on the mRNA expression of the endogenous PR gene. The PR is a well-known ER regulated gene, and its expression in breast cancer cells is used as a marker for functional ER signalling (Clarke *et al.* 1994, Yang *et al.* 2001, Rickard *et al.* 2002, Han *et al.* 2008). Furthermore, it has previously been shown that E₂-activated ER β , like ER α , enhances the expression of the endogenous PR gene in the uterine and bone cells (Schultz *et al.* 2005). Surprisingly, although we showed that the AR, both unliganded and liganded, had no effect on ER β -mediated transactivation in an ERE-reporter assay (figure 3.5B and 3.7A-C), the results in figure 3.8C showed that the unliganded AR downregulates the ER β -mediated regulation of the endogenous PR gene. In contrast, the ability of the unliganded AR to inhibit ER α -mediated transactivation on the synthetic promoter-reporter gene (figure 3.2B) was replicated on the endogenous PR gene (figure 3.8B) with a similar percentage reduction observed at the higher AR concentrations (2- and 4-fold). These results indicate that although effects seen on a synthetic promoter-reporter

can often be reproduced on an endogenous gene, this is not always the case. Effects on the synthetic promoter are likely due to the AR and ER α competing for binding to ERE, as previously reported by Peters *et al.* (2009). However, considering that the AR inhibits the endogenous PR gene in the presence of either ER α or ER β , it may be that effects on the endogenous promoter is more complex than effects on a simple synthetic ERE-promoter.

Although the general mechanism for transactivation is via the interaction of the activated ER with EREs in target genes, the human PR does not contain a palindromic ERE (Misrahi *et al.* 1988, Kastner *et al.* 1990, Schultz *et al.* 2005). Instead, it contains an ERE half-site and two specificity protein (Sp1) sites, all of which were previously shown to be needed for E₂-mediated activation of the PR gene (Schultz *et al.* 2003, Petz *et al.* 2004). Thus it may mean that ER α - and ER β -mediated activation of the endogenous PR gene by E₂, requires both the ERE half site and the Sp1 sites. The possibility that the AR competes with either ER α or ER β for the ERE half-site, can thus not be excluded. Furthermore, as the AR has been shown to repress Sp1-induced transcription (Curtin *et al.* 2001, Verras *et al.* 2007, Grosse *et al.* 2012), it may be possible for the AR to compete with both ER α and ER β for tethering to Sp1, thereby inhibiting the activation of the PR gene. Taken together, we suggest a putative mechanism whereby the AR and ER α , but not ER β , compete for binding to the synthetic ERE, while on the endogenous PR gene both ER α and ER β compete with the AR for binding to either the ERE half-sites and/or for tethering to Sp1. However, further studies are needed to support this hypothesis.

4.4. E₂ upregulation of the endogenous ARE-containing PSA gene can be mediated by either ER α or ER β

As mentioned previously, Peters and co-workers showed that the AR can bind to either an ARE or ERE, while ER α only binds to an ERE (Peters *et al.* 2009). However, we showed that E₂ via ER α could transactivate a simple ARE-containing promoter (figure 3.9B), as well as

upregulating the mRNA expression of the endogenous ARE-containing PSA gene in MDA-MB-231 cells (figure 3.10B). These results suggest, at least for the simple ARE in the current experimental system that ER α binds to the ARE. In addition, we show that E₂ upregulated PSA mRNA expression in the presence of overexpressed ER β (figure 3.10A) but did not transactivate the simple ARE via ER β (figure 3.9A), yet again, indicating that effects seen on a synthetic promoter cannot always be reproduced on an endogenous gene, and that regulation of an endogenous gene is more complex than regulation of a simple synthetic promoter. For the endogenous PSA gene, the possibility that ER α and ER β could be interacting with *cis*-elements other than the ARE in the PSA promoter cannot be excluded. It is known that Sp1 binding sites are found in the PSA promoter (Sadar *et al.* 1999, Kim and Coetzee 2004), and as previously mentioned, it is known that ER α can tether to the Sp1 transcription factor to activate the transcription of the Sp1 containing PR gene (Petz *et al.* 2004, Schultz *et al.* 2005). Surprisingly, a 6.0-fold increase in the expression of the PSA gene was observed in the presence of ER β and E₂ (figure 3.10A), while only a 2.1-fold was seen in the presence of the AR and DHT (figure 3.10C). However, it should be noted that the latter was the result of a single experiment, and should be repeated before any conclusions are made.

Interestingly the results in figure 3.9B showed that DHT also activates the simple ARE-containing promoter via ER α (a 3.7-fold increase). Although it appears that a similar activation (1.8-fold) of the endogenous PSA gene occurs (figure 3.10B), it was not statistically significant. To the best of our knowledge, there are no reports in the literature on the activation of ER α by DHT. However, at least one study has reported that DHT binds weakly to rat ER α and ER β (Kuiper *et al.* 1997).

In summary, our results in MDA-MB-231 cells indicating that ER α can transactivate an ARE-containing promoter/gene are in contrast to the findings reported by (Peters *et al.* 2009) showing that ER α cannot bind to AREs in COS-1 and T47D cells. Taking into consideration that

receptors and their hormones behave differently in different cell lines (Birrell *et al.* 1998), the discrepancy in results could be due to the different techniques and/or cell lines used.

4.5. The unliganded AR impedes repression of transcription via ER α , but not ER β , while the liganded AR impedes the transrepression function of both ER α and ER β .

Most studies on the ER focus on the mechanism by which ERs regulate the transcription of E₂ regulated genes via direct binding to EREs. As described in Chapter 1 (Section 1.3.2.2), the ERs can also repress transcription of target genes by tethering to other DNA-bound transcription factors such as NF κ B (Stein and Yang 1995, Galien and Garcia 1997). Thus, we next investigated whether the AR could modulate repression of transcription via the ER subtypes. In agreement with previously reported studies (Stein and Yang 1995, Galien and Garcia 1997, Visser *et al.* 2013), we showed that E₂ repressed the PMA-induced activation of a simple NF κ B-containing promoter-reporter construct via both ER α and ER β (figure 3.11B and D).

Similar to the results for transactivation of a simple ERE-containing promoter-reporter construct showing that the unliganded AR inhibits the transactivation function of ER α but not of ER β (figure 3.2B and 3.5B), the results in figure 3.11B and 3.11D show that the unliganded AR inhibits the E₂-induced repression of NF κ B activity via ER α but not via ER β . To the best of our knowledge, this study is the first to show that the AR can modulate the transrepressive effects of E₂ via ER α . It is noteworthy that this modulation by the AR is not seen for ER β . The inhibition noted in the case of ER α may be due to the unliganded and liganded AR interacting with ER α , physically preventing ER α from tethering to NF κ B.

As seen for transactivation, the AR ligands do not modulate AR effects on ER α -mediated transrepression. Surprisingly, the results in figure 3.12A-C show that ligand-bound AR was able to inhibit the transrepression function of ER β . Additional studies are needed to elucidate the mechanism behind this finding, as well as to evaluate its physiological impact. Considering that both the AR and ER subtypes can tether to NF κ B leading to repression (Palvimo *et al.* 1996, Aarnisalo *et al.* 1998, McKay and Cidlowski 1999), it may be possible that the AR and ER subtypes are competing for binding to NF κ B, thus the activated AR may prevent ER β from causing transrepression thereby inhibiting the ER-mediated anti-inflammatory effects of E₂. It is known that inflammation plays an important role in breast cancer development and progression (Niu *et al.* 2012, Bhatelia *et al.* 2014, Suman *et al.*), thus our results showing that the AR reduces the anti-inflammatory effects of the ER may suggest that inflammation-induced breast cancer progression is promoted. However further studies are needed to assess the link between the ER, inflammation and breast cancer.

4.6. No change in proliferation was observed in cells transfected with ER

Transactivation and transrepression of ER target genes aid in the regulation of breast cancer cell proliferation (McKay and Cidlowski 1999, Beato and Klug 2000). Since the results from the current study showed that the unliganded AR could inhibit the transactivation and transrepression function of ER α , but not of ER β , using simple ERE- and NF κ B-containing promoter reporter constructs, we next evaluated the effects of the AR on E₂-stimulated proliferation of the human MDA-MB-231 breast cancer cell line. A MTT cell proliferation assay was thus performed in MDA-MB-231 cells transfected with either ER α or ER β in the absence of the AR. As we did not detect E₂-induced proliferation in either the untransfected MDA-MB-231 cells or in those with overexpressed ER α or ER β (figure 3.14), we could not evaluate the modulatory effects of the AR on cell proliferation.

Upon closer inspection of the literature, it appears that most studies employ the ER-positive MCF-7 breast cancer cell line (Han *et al.* 2008, Zhang *et al.* 2013) when investigating E₂-mediated cell proliferation. This is probably due to the fact that a number of studies have shown that cell proliferation is inhibited when ER α or ER β is reintroduced into MDA-MB-231 cells (Garcia *et al.* 1992, Levenson and Jordan 1994, Lazennec *et al.* 2001, Moggs *et al.* 2005).

A possible explanation for the discrepancy in the results in the current study showing no proliferation versus the above-mentioned studies showing inhibition of proliferation, may be due to different experimental conditions used. For example, Garcia *et al.* (1992) treated MDA-MB-231 cells for 9 days and Lazennec *et al.* (2001) treated for 6 days, while in the current study cells were incubated for only 2 days. Furthermore, understanding why E₂ has an inhibitory effect on the proliferation of ER-negative cells when the ER is re-introduced, is a topic under investigation by a number of researchers (Garcia *et al.* 1992, Lazennec *et al.* 2001, Moggs *et al.* 2005).

4.7. Conclusions and Future Work

Several studies have shown that E₂ and ER play critical roles in breast cancer biology and their role in breast cancer has been extensively studied (Leygue *et al.* 1998, Speirs *et al.* 1999, Helguero *et al.* 2005, Yager and Davidson 2006, Zhao *et al.* 2008, Hartman *et al.* 2009). The ER α has been reported to promote cell growth and survival, while ER β inhibits the ER α -mediated cell proliferation (Leygue *et al.* 1998, Dotzlaw *et al.* 1999, Speirs *et al.* 1999, Zhao *et al.* 2008). The role of other steroid receptors such as the AR in breast cancer is, however, an area of ongoing research, with emerging evidence suggesting that the AR is also involved in the regulation of breast cancer growth (Birrell *et al.* 1998, Brys 2000, Hartman *et al.* 2012, Shah *et al.* 2013, Lakis *et al.* 2014). For example, previous evidence suggests that the AR has the ability to inhibit the transactivation function of ER α , with not much known about the crosstalk

between the AR and the ER β . Considering that the presence of either the AR or ER β in ER α -positive breast cancers has been shown to be beneficial (Birrell *et al.* 1995, Szelei *et al.* 1997, Lanzino *et al.* 2005, Macedo *et al.* 2006, Williams *et al.* 2007, Hartman *et al.* 2009, Peters *et al.* 2009, Hickey *et al.* 2012), and that there is a lack of research investigating the interplay between the AR and ER β in breast cancer, the aim of the current study was to elucidate the effects of the AR on the activity of ER β .

The results presented in Chapter 3, showed that although the AR has no effect on the transactivation function of ER β on a simple, synthetic ERE-containing promoter, it downregulates the ER β -driven mRNA expression of the endogenous PR gene, a well-described ER-regulated gene. Considering that upregulation of this ER-regulated gene is associated with an increased risk of breast cancer (Fox *et al.* 2008, Han *et al.* 2008, Sivik and Jansson 2012) and evidence shows that ER β acts like ER α when expressed in breast cancers lacking ER α , our results suggest that the AR may have a protective role in breast cancers expressing either ER β or ER α . However, in inflammation-induced breast cancer the AR may not be protective considering results obtained in the current study for transrepression. Specifically, for transrepression studies, we show for the first time that both the unliganded- and ligand-bound AR inhibits the transrepression function of ER α , while only the ligand-bound AR was able to inhibit the transrepression function of ER β . As it is known that inflammation plays a role in the development and progression of breast cancer (Niu *et al.* 2012, Bhatelia *et al.* 2014, Suman *et al.* 2015), our results suggest that the unliganded- or ligand-bound AR may promote breast cancer risk by reducing the anti-inflammatory effects of ER α , while the AR can only do so for ER β when ligand bound. Furthermore, the current study also shows ER α -mediated, E₂-induced transactivation via a synthetic ARE as well as via an endogenous ARE-containing promoter, while the same is true for ER β only via an endogenous ARE-containing promoter. It is interesting to note that ER β seems to be upregulating the transcription of the endogenous PR and PSA genes via a mechanism involving tethering of ER β to the Sp1 transcription factor.

While we cannot exclude that ER α also acts via tethering to Sp1, it appears that ER α activates transcription primarily via binding to EREs.

To the best of our knowledge this study is the first to show that the activated AR affects the transrepression function of both ER α and ER β on a synthetic NF κ B-containing promoter. A limitation of this study is that we did not investigate the influence of the AR on ER-mediated transrepression of any endogenous NF κ B-containing promoters. This could be addressed on endogenous genes such as interleukin (IL)-6 or Regulated on Activation, Normal T-cell Expressed, and Secreted (RANTES) (Luboshits *et al.* 1999, Knüpfer and Preiß 2007), which are known to be involved in the development and progression of breast cancer and contain NF κ B-binding sites within their promoters (Libermann and Baltimore 1990, Li and Bever 2001, Azenshtein *et al.* 2002). To investigate whether the presence of the AR inhibits the ability of the ER subtypes to tether to NF κ B, as suggested in Section 4.5, co-immunoprecipitation (Co-IP) and/or fluorescence resonance energy transfer (FRET) assays may be used. Co-IP assays are used to investigate protein-protein interactions *in vitro* (Fields and Song 1989), while the FRET assay allows the investigation of protein-protein interaction in living cells (Clegg 1995). A further limitation was that we did not plate into a 10 cm dish for transfections, therefore protein concentration was not the best normalization method.

Lastly, as no E₂-stimulated proliferation could be obtained in the ER-negative breast cancer cell line containing overexpressed ER α or ER β , the effect of the AR on ER-mediated cell proliferation could not be investigated. Proliferation assays should thus be carried out in a cell line which is known to express both ER subtypes such as the MCF-7 or T47D breast cancer cell line. The role of the AR on ER α - or ER β -mediated breast cancer cell proliferation can be determined in these cell lines by silencing the expression of the respective ER subtype using siRNA technology

To gain more insight into how the AR modulates ER β -mediated regulation of the PR gene, chromatin immunoprecipitation (ChIP) assays can be used to firstly determine whether ER β is recruited to *cis*-elements such as the ERE-half sites and/or Sp1 binding sites in the PR promoter. Secondly, ChIP can be used to determine whether the AR competes with ER β for binding to the above-mentioned *cis*-regulatory elements as it has previously been shown that the AR can bind to the ERE and can tether to the Sp1 (Curtin *et al.* 2001, Verras *et al.* 2007, Peters *et al.* 2009, Grosse *et al.* 2012). The ChIP assay is considered a relatively inexpensive method to study the interaction between a protein and a specific DNA sequence *in vivo*. Furthermore, to establish whether the AR interacts with ER β and the Sp1 transcription factor to form a complex on the promoter of the PR gene, sequential chromatin immunoprecipitation (re-ChIP) assays may be employed. The mechanism whereby ER α and ER β upregulates the mRNA expression of the PSA gene, which contains AREs, Sp1 binding sites as well as other *cis* regulatory elements, can also be further investigated by ChIP assays. Besides studying the recruitment of the ER subtypes to the Sp1 *cis*-regulatory element, similar experiments as described above may be used to determine whether the ER subtypes interact with the Sp1 transcription factor on the PSA promoter.

To examine whether the Sp1 transcription factor is required for ER β -mediated upregulation of PR gene expression, siRNA technology can be used to silence the expression of Sp1. In addition, it would be interesting to examine the effects of the AR on other ER β -regulated endogenous genes such as cathepsin D (CTSD) (Behrens *et al.* 2007), as it has previously been shown that the AR inhibits the expression of the ER regulated PR gene, but not the CTSD gene (Peters *et al.* 2009). Lastly, although the use of a yeast two-hybrid assay has previously shown that the AR and ER β do not interact with each other, it would be of interest to determine whether a direct association between these steroid receptors exists in our model system. The latter could be investigated using either Co-IP or FRET assays.

In summary, the results presented in this thesis showing that the AR inhibits the activity of ER α via a synthetic ERE-promoter as well as an endogenous ER-regulated gene and that of ER β via an endogenous ER-regulated gene, may suggest that the presence of AR is also a good prognostic marker for breast cancer patients as previously reported for ER α by Peters *et al.* (2009). On the other hand, our novel results showing that the activated AR reduces the anti-inflammatory effects of the ER, may suggest that the AR promotes inflammatory-induced breast cancer. The precise physiological implications of these results remain to be determined. Finally, although the results from this study are preliminary and have certain limitations, it nonetheless highlights the fact that the role of the AR in breast cancer is a complex one. Our findings may thus aid in understanding of crosstalk between the ER and AR signalling pathways, and how it contributes to the growth and survival of breast cancer cells. Figure 4.1 shows the contribution of findings of this current study to the existing knowledge on the interplay between the AR and ER β .




Figure 4.1. Interplay between the ER subtypes and the AR may lead to both positive and negative effects on breast cancer. ERα promotes cell survival and proliferation, while ERβ antagonizes ERα-mediated cell proliferation. The AR has been shown to have the ability to inhibit cell proliferation both directly and indirectly by suppressing the activity of ERα. The AR also has no effect on the activity of ERβ on synthetic ERE-containing promoter, while the AR inhibits the upregulation of the ER-regulated PR gene. Lastly the AR promotes inflammation-induced breast cancer by inhibiting the transrepression function of both ERα and ERβ. – and T denote inhibition of activity, + shows activation of activity, the yellow circle with a line inside indicates no effect on the activity, the blue arrows demonstrate that the effect is towards ERα while the purple arrow demonstrates that the effect is towards ERβ.

REFERENCES

AARNISALO, P., J. J. PALVIMO and O. A. JÄNNE (1998). "CREB-binding protein in androgen receptor-mediated signaling." Proceedings of the National Academy of Sciences **95**(5): 2122-2127.

ABDULJABBAR, R., O. NEGM, C.-F. LAI, D. JERJEES, M. AL-KAABI, M. HAMED, P. TIGHE, L. BULUWELA, A. MUKHERJEE, A. GREEN, S. ALI, E. RAKHA and I. ELLIS (2015). "Clinical and biological significance of glucocorticoid receptor (GR) expression in breast cancer." Breast Cancer Research and Treatment **150**(2): 335-346.

ABPLANALP, W., M. RYMASZEWSKI, J. ADAMSKI and M. T. R. SUBBIAH (1999). "Evidence for interference in estradiol-17 β inactivation to estrone by oxidized low-density lipoprotein and selected lipid peroxidation products." Journal of Laboratory and Clinical Medicine **134**(3): 253-259.

AFRICANDER, D., K.-H. STORBECK and J. HAPGOOD (2014). "A comparative study of the androgenic properties of progesterone and the progestins, medroxyprogesterone acetate (MPA) and norethisterone acetate (NET-A)." Journal of Steroid Biochemistry and Molecular Biology **143**: 404-415.

AFRICANDER, D., N. VERHOOG and P. HAPGOOD, JANET. (2011). "Molecular mechanisms of steroid receptor-mediated actions by synthetic progestins used in HRT and contraception." Steroids **76**: 636-652.

AKÇAY, H., TÜRKER. and R. BAYRAK (2014). "Computational studies on the anastrozole and letrozole, effective chemotherapy drugs against breast cancer." Spectrochimica Acta Part A: Molecular and Biomolecular Spectroscopy **122**(0): 142-152.

ALARID, E. T., B. NATALIE and S. NATALIA (1999). "Proteasome-Mediated Proteolysis of Estrogen Receptor: A Novel Component in Autologous Down-Regulation." Molecular Endocrinology **13**(9): 1522-1534.

ALSHENAWY, H. A. (2012). "Prevalence of androgen receptors in invasive breast carcinoma and its relation with estrogen receptor, progesterone receptor and Her2/neu expression." Journal of the Egyptian National Cancer Institute **24**(2): 77-83.

ANBALAGAN, M., B. HUDERSON, L. MURPHY and B. G. ROWAN (2012). "Post-translational modifications of nuclear receptors and human disease." Nuclear receptor signaling **10**.

ANDERSSON, S. and N. MOGHRABI (1997). "Physiology and molecular genetics of 17 β -hydroxysteroid dehydrogenases." Steroids **62**(1): 143-147.

ANDO, S., F. DE AMICIS, V. RAGO, A. CARPINO, M. MAGGIOLINI, M. L. PANNO and M. LANZINO (2002). "Breast cancer: from estrogen to androgen receptor." Molecular and cellular endocrinology **193**(1): 121-128.

ANESTIS, A., M. V. KARAMOUZIS, G. DALAGIORGOU and A. G. PAPAVASSILIOU (2015). "Is androgen receptor targeting an emerging treatment strategy for triple negative breast cancer?" Cancer Treatment Reviews(0).

ANTOINE, C., L. AMEYE, M. PAESMANS and S. ROZENBERG (2014). "Systematic review about breast cancer incidence in relation to hormone replacement therapy use." Climacteric **17**(2): 116-132.

ASCENZI, P., A. BOCEDI and M. MARINO (2006). "Structure–function relationship of estrogen receptor α and β : Impact on human health." Molecular Aspects of Medicine **27**(4): 299-402.

ASCH-KENDRICK, R. J., M. A. SAMOLS, M. T. LILO, A. P. SUBHAWONG, R. SHARMA, P. B. ILLEI, P. ARGANI and A. CIMINO-MATHEWS (2014). "NKX3.1 is expressed in ER-positive and AR-positive primary breast carcinomas." Journal of clinical pathology **67**(9): 768-771.

ASSIKIS, V. J. and J. CRAIG, V. (1997). "Risks and benefits of tamoxifen therapy." Oncology **11**(2 SUPPL. 1): 21-23.

AZENSHTAIN, E., G. LUBOSHITS, S. SHINA, E. NEUMARK, D. SHAHBAZIAN, M. WEIL, N. WIGLER, I. KEYDAR and A. BEN-BARUCH (2002). "The CC Chemokine RANTES in Breast Carcinoma Progression: Regulation of Expression and Potential Mechanisms of Promalignant Activity." Cancer Research **62**(4): 1093-1102.

BAIN, D., L., A. HENEGHAN, F., K. CONNAGHAN-JONES, D. and M. MIURA, T. (2007). "Nuclear Receptor Structure: Implications for Function." Annual Review of Physiology **69**(1): 201-220.

BALFE, P. J., A. H. MCCANN, H. M. WELCH and M. J. KERIN (2004). "Estrogen receptor β and breast cancer." European Journal of Surgical Oncology (EJSO) **30**(10): 1043-1050.

BEATO, M., G. CHALEPAKIS, M. SCHAUER and E. P. SLATER (1989). "DNA regulatory elements for steroid hormones." Journal of Steroid Biochemistry **32**(5): 737-747.

BEATO, M. and J. KLUG (2000). "Steroid hormone receptors: an update." Human Reproduction Update **6**(3): 225-236.

BECK, I., M, E., V. B. WIM, V. LINDA, R. Y. KEITH, H. GUY and D. B. KAROLIEN (2009). "Crosstalk in Inflammation: The Interplay of Glucocorticoid Receptor-Based Mechanisms and Kinases and Phosphatases." Endocrine Reviews **30**(7): 830-882.

BEHRENS, D., J. H. GILL and I. FICHTNER (2007). "Loss of tumorigenicity of stably ER β -transfected MCF-7 breast cancer cells." Molecular and Cellular Endocrinology **274**(1-2): 19-29.

BERAL, V. (2003). "Breast cancer and hormone-replacement therapy in the Million Women Study." The Lancet **362**(9382): 419-427.

BERGKVIST, L., H.-O. ADAMI, I. PERSSON, R. HOOVER and C. SCHAIRER (1989). "The Risk of Breast Cancer after Estrogen and Estrogen-Progestin Replacement." New England Journal of Medicine **321**(5): 293-297.

BERGMANN, A., C. EULENBERG, M. WELLNER, S. ROLLE, F. LUFT and R. KETTRITZ (2010). "Aldosterone Abrogates Nuclear Factor κ B-Mediated Tumor Necrosis Factor α Production in Human Neutrophils via the Mineralocorticoid Receptor." Hypertension **55**(2): 370-379.

BHASIN, S. and R. JASUJA (2009). "Selective Androgen Receptor Modulators (SARMs) as Function Promoting Therapies." Current opinion in clinical nutrition and metabolic care **12**(3): 232-240.

BHATELIA, K., K. SINGH and R. SINGH (2014). "TLRs: Linking inflammation and breast cancer." Cellular Signalling **26**(11): 2350-2357.

BIRRELL, N., STEPHEN., HALL R.E and W. TILLEY (1998). "Role of the Androgen Receptor in Human Breast Cancer." Journal of Mammary Gland Biology and Neoplasia **3**(1): 95-103.

BIRRELL, S. N., J. M. BENTEL, T. E. HICKEY, C. RICCIARDELLI, M. A. WEGER, D. J. HORSFALL and W. D. TILLEY (1995). "Androgens induce divergent proliferative responses in human breast cancer cell lines." The Journal of Steroid Biochemistry and Molecular Biology **52**(5): 459-467.

BIRRELL, S. N., L. M. BUTLER, J. M. HARRIS, G. BUCHANAN and W. D. TILLEY (2007). "Disruption of androgen receptor signaling by synthetic progestins may increase risk of developing breast cancer." The FASEB Journal **21**(10): 2285-2293.

BITZER, J., P. KENEMANS and A. MUECK, O. (2008). "Breast cancer risk in postmenopausal women using testosterone in combination with hormone replacement therapy." Maturitas **59**(3): 209-218.

BJÖRNSTRÖM, L. and M. SJÖBERG (2005). "Mechanisms of Estrogen Receptor Signaling: Convergence of Genomic and Nongenomic Actions on Target Genes." Molecular Endocrinology **19**(4): 833-842.

BLACK, D. M. (1994). "Special Issue: Genetics of CancerThe genetics of breast cancer." European Journal of Cancer **30**(13): 1957-1961.

BLACK, M. H., M. GIAI, R. PONZONE, P. SISMONDI, H. YU and E. P. DIAMANDIS (2000). "Serum total and free prostate-specific antigen for breast cancer diagnosis in women." Clinical cancer research **6**(2): 467-473.

BOHRER, R. L., SHUAI. CHEN, TIMOTHY. C. HALLSTROM and HAOJIE. HUANG (2010). "Androgens Suppress EZH2 Expression Via Retinoblastoma (RB) and p130-Dependent Pathways: A Potential Mechanism of Androgen-Refractory Progression of Prostate Cancer." Endocrinology **151**(11): 5136-5145.

BOONYARATANAKORNKIT, V., Y. BI, M. RUDD and D. EDWARDS, P. (2008). "The role and mechanism of progesterone receptor activation of extra-nuclear signaling pathways in regulating gene transcription and cell cycle progression." Steroids **73**(9–10): 922-928.

BOTELHO, M., C., H. ALVES, A. BARROS, G. RINALDI, P. J. BRINDLEY and M. SOUSA (2015). "The role of estrogens and estrogen receptor signaling pathways in cancer and infertility: the case of schistosomes." Trends in Parasitology(0).

BRADFORD, M. (1976). "A Rapid and Sensitive Method for the Quantitation of Microgram Quantities of Protein Utilizing the Principle of Protein-Dye Binding " Analytical Biochemistry **72**: 248-254.

BRANDENBERGER, A., W. , T. MENG KIAN , J. LEE, Y. , V. CHAO and R. JAFFE, B. (1997). "Tissue Distribution of Estrogen Receptors Alpha (ER- α) and Beta (ER- β) mRNA in the Midgestational Human Fetus." The Journal of Clinical Endocrinology & Metabolism **82**(10): 3509-3512.

BRENTANI, M., MITZI, E. FRANCO, LUIS., C. OSHIMA, TIZUKO, F. and M. PACHECO, MEDEIROS. (1986). "Androgen, estrogen, and progesterone receptor levels in malignant and benign breast tumors: A multivariate analysis approach." International Journal of Cancer **38**(5): 637-642.

BRINKMANN, U., R. MATTES, E. and P. BUCKEL (1989). "High-level expression of recombinant genes in Escherichia coli is dependent on the availability of the dnaY gene product." Gene **85**(1): 109-114.

BRISKEN, C. and B. O'MALLEY (2010). "Hormone action in the mammary gland." Cold Spring Harbor perspectives in biology **2**(12): a003178.

- BROSENS, J., J., J. TULLET, R. VARSHOCHI and E. LAM, W. F. (2004). "Steroid receptor action." Best Practice & Research Clinical Obstetrics & Gynaecology **18**(2): 265-283.
- BROWN, S., B. and S. HANKINSON, E. (2014). "Endogenous estrogens and the risk of breast, endometrial, and ovarian cancers." Steroids(0).
- BRYN, M. (2000). "Androgens and the androgen receptor: do they play a role in breast cancer?" Med Sci Monit **6**(2): 433-438.
- BUCHANAN, G., S. BIRRELL, N., A. PETERS, A., T. BIANCO-MIOTTO, K. RAMSAY, E. COPS, J., M. YANG, J. HARRIS, M., H. A. SIMILA and N. L. MOORE (2005). "Decreased androgen receptor levels and receptor function in breast cancer contribute to the failure of response to medroxyprogesterone acetate." Cancer research **65**(18): 8487-8496.
- BUSTIN, S. A. (2002). "Quantification of mRNA using real-time reverse transcription PCR (RT-PCR): Trends and problems." Journal of Molecular Endocrinology **29**(1): 23-39.
- CARTER, L., ERIC. and W. STEPHEN, RAGSDALE. (2014). "Modulation of nuclear receptor function by cellular redox poise." Journal of Inorganic Biochemistry **133**: 92-103.
- CASTELLANO, I., A. ELENA, A. VALERIA, V. ANNA MARIA, C. LUIGI, A. RICCARDO, D. ANTONIO, D. MICHELA, B. GIANNI, S. ALAN, COATES., V. GIUSEPPE and S. ANNA (2010). "Androgen receptor expression is a significant prognostic factor in estrogen receptor positive breast cancers." Breast cancer research **124**: 607-617.
- CAULEY, J. A., F. L. LUCAS, L. H. KULLER, K. STONE, W. BROWNER and S. R. CUMMINGS (1999). "Elevated Serum Estradiol and Testosterone Concentrations Are Associated with a High Risk for Breast Cancer." Annals of Internal Medicine **130**(4_Part_1): 270-277.
- CELI, F., S., M. COHEN, M., S. ANTONARAKIS, E., E. WERTHEIMER, J. ROTH and A. SHULDINER, R. (1994). "Determination of Gene Dosage by a Quantitative Adaptation of the Polymerase Chain Reaction (gd-PCR): Rapid Detection of Deletions and Duplications of Gene Sequences." Genomics **21**(2): 304-310.
- CHANG, M.-S., B.-C. CHEN, M.-T. YU, J.-R. SHEU, T.-F. CHEN and C.-H. LIN (2005). "Phorbol 12-myristate 13-acetate upregulates cyclooxygenase-2 expression in human pulmonary epithelial cells via Ras, Raf-1, ERK, and NF- κ B, but not p38 MAPK, pathways." Cellular Signalling **17**(3): 299-310.
- CHANPLAKORN, N., P. CHANPLAKORN, T. SUZUKI, K. ONO, L. WANG, M. S. CHAN, L. WING, C. C. YIU, L. W.-C. CHOW and H. SASANO (2011). "Increased 5 α -reductase type 2 expression in human breast carcinoma following aromatase inhibitor therapy: the correlation with decreased tumor cell proliferation." Hormones and Cancer **2**(1): 73-81.

CHANTONG, B., D. KRATSCHMAR, V., L. NASHEV, G., Z. BALAZS and A. ODERMATT (2012). "Mineralocorticoid and glucocorticoid receptors differentially regulate NF-kappaB activity and pro-inflammatory cytokine production in murine BV-2 microglial cells." Journal of Neuroinflammation **9**: 260-260.

CHARITIDI, K., I. MELTSEY, Y. TAHERA and B. CANLON (2009). "Functional responses of estrogen receptors in the male and female auditory system." Hearing Research **252**(1-2): 71-78.

CHATTERTON, J., T. ROBERT, A. GEIGER, S., P. GANN, H. and S. KHAN, A. (2003). "Formation of estrone and estradiol from estrone sulfate by normal breast parenchymal tissue." The Journal of Steroid Biochemistry and Molecular Biology **86**(2): 159-166.

CHEN, Y.-J., C.-M. HUNG, N. KAY, C.-C. CHEN, Y.-H. KAO and S.-S. YUAN (2012). "Progesterone receptor is involved in 2,3,7,8-tetrachlorodibenzo-p-dioxin-stimulated breast cancer cells proliferation." Cancer Letters **319**(2): 223-231.

CHLEBOWSKI, R. T., G. L. ANDERSON and M. GASS (2010). "Estrogen plus progestin and breast cancer incidence and mortality in postmenopausal women." JAMA **304**(15): 1684-1692.

CHLEBOWSKI, R. T., S. L. HENDRIX and R. D. LANGER (2003). "Influence of estrogen plus progestin on breast cancer and mammography in healthy postmenopausal women: The women's health initiative randomized trial." JAMA **289**(24): 3243-3253.

CHOTTANAPUND, S., M. B. M. VAN DUURSEN, P. NAVASUMRIT, P. HUNSONTI, S. TIMTAVORN, M. RUCHIRAWAT and M. VAN DEN BERG (2013). "Effect of androgens on different breast cancer cells co-cultured with or without breast adipose fibroblasts." The Journal of Steroid Biochemistry and Molecular Biology **138**(0): 54-62.

CIOCCA, D., R., F. GAGO, E., M. FANELLI, L. A. and S. CALDERWOOD, K. (2006). "Co-expression of steroid receptors (estrogen receptor alpha and/or progesterone receptors) and Her-2/neu: Clinical implications." The Journal of Steroid Biochemistry and Molecular Biology **102**(1-5): 32-40.

CIRUELOS, E., T. PASCUAL, M. L. ARROYO VOZMEDIANO, M. BLANCO, L. MANSO, L. PARRILLA, C. MUÑOZ, E. VEGA, M. J. CALDERÓN, B. SANCHE and H. CORTES-FUNES (2014). "The therapeutic role of fulvestrant in the management of patients with hormone receptor-positive breast cancer." The Breast **23**(3): 201-208.

CLARKE, C. L. and R. L. SUTHERLAND (1990). "Progestin Regulation of Cellular Proliferation." Endocrine Reviews **11**(2): 266-301.

CLARKE, R., C. MINETTA, LIU., B. KERRIE, BOUKER., G. ZHIPING, Y. RICHARD, LEE., Z. YUELIN, C. TODD, SKAAR., G. BIANCA, O. B. KERRY, W. YUE and A. LEENA,

HILAKIVI-CLARKE. (2003). "Antiestrogen resistance in breast cancer and the role of estrogen receptor signaling." Oncogene **22**: 7316-7339.

CLARKE, R., T. SKAAR, K. BAUMANN, F. LEONESSA, M. JAMES, J. LIPPMAN, E. THOMPSON, C. FRETER and N. BRUNNER (1994). "Hormonal carcinogenesis in breast cancer: cellular and molecular studies of malignant progression." Breast Cancer Research and Treatment **31**(2-3): 237-248.

CLARKE, R. B., A. HOWELL, C. S. POTTEN and E. ANDERSON (1997). "Dissociation between steroid receptor expression and cell proliferation in the human breast." Cancer research **57**(22): 4987-4991.

CLEGG, R. M. (1995). "Fluorescence resonance energy transfer." Current Opinion in Biotechnology **6**(1): 103-110.

COCHRANE, D. R., S. BERNALES, B. M. JACOBSEN, D. M. CITTELLY, E. N. HOWE, N. C. D'AMATO, N. S. SPOELSTRA, S. M. EDGERTON, A. JEAN and J. GUERRERO (2014). "Role of the androgen receptor in breast cancer and preclinical analysis of enzalutamide." Breast Cancer Res **16**(1): R7.

COLDITZ, G. A., S. E. HANKINSON, D. J. HUNTER, W. C. WILLETT, J. E. MANSON, M. J. STAMPFER, C. HENNEKENS, B. ROSNER and F. E. SPEIZER (1995). "The use of estrogens and progestins and the risk of breast cancer in postmenopausal women." New England Journal of Medicine **332**(24): 1589-1593.

CONNELLY, O., M., B. JERICEVIC, M. and J. LYDON, P. (2003). "Progesterone Receptors in Mammary Gland Development and Tumorigenesis." Journal of Mammary Gland Biology and Neoplasia **8**(2): 205-214.

CONZEN, S., D. (2008). "Minireview: Nuclear Receptors and Breast Cancer." Molecular Endocrinology **22**(10): 2215-2228.

CORTHALS, G. L., B. LIN, A. MONSON, V. NGUYEN, B. R. FRANZA, S. R. PLYMATE, R. AEBERSOLD and L. HOOD (2000). "Comprehensive analyses of prostate gene expression: convergence of expressed sequence tag databases, transcript profiling and proteomics." Electrophoresis **21**: 1823-1831.

COS, S., R. FERNÁNDEZ, A. GÜÉZMES and E. J. SÁNCHEZ-BARCELÓ (1998). "Influence of Melatonin on Invasive and Metastatic Properties of MCF-7 Human Breast Cancer Cells." Cancer Research **58**(19): 4383-4390.

COSS, C., C., A. JONES and J. DALTON, T. (2014). "Selective androgen receptor modulators as improved androgen therapy for advanced breast cancer." Steroids **90**(0): 94-100.

COUSE, J. F. and K. S. KORACH (1999). "Estrogen receptor null mice: What have we learned and where will they lead us?" Endocrine Reviews **20**(3): 358-417.

CUNAT, S., P. HOFFMANN and P. PUJOL (2004). "Estrogens and epithelial ovarian cancer." Gynecologic Oncology **94**(1): 25-32.

CUNHA, G., P. YOUNG, Y. HOM, P. COOKE, J. TAYLOR and D. LUBAHN (1997). "Elucidation of a role for stromal steroid hormone receptors in mammary gland growth and development using tissue recombinants." Journal of mammary gland biology and neoplasia **2**(4): 393-402.

CURTIN, D., S. JENKINS, N. FARMER, A. C. ANDERSON, D. J. HAISENLEDER, E. RISSMAN, E. M. WILSON and M. A. SHUPNIK (2001). "Androgen suppression of GnRH-stimulated rat LH β gene transcription occurs through Sp1 sites in the distal GnRH-responsive promoter region." Molecular Endocrinology **15**(11): 1906-1917.

CVORO, A., C. YUAN, S. PARUTHIYIL, O. MILLER, H., K. YAMAMOTO, R. and D. LEITMAN, C. (2011). "Cross Talk between Glucocorticoid and Estrogen Receptors Occurs at a Subset of Proinflammatory Genes." The Journal of Immunology **186**(7): 4354-4360.

DAUVOIS, S., C.-S. GENG, C. LÉVESQUE, Y. MÉRAND and F. LABRIE (1991). "Additive Inhibitory Effects of an Androgen and the Antiestrogen EM-170 on Estradiol-stimulated Growth of Human ZR-75-1 Breast Tumors in Athymic Mice." Cancer Research **51**(12): 3131-3135.

DAUVOIS, S. and F. LABRIE (1989). "Androstenedione and androst-5-ene-3 β , 17 β -diol stimulate DMBA-induced rat mammary tumors—role of aromatase." Breast cancer research and treatment **13**(1): 61-69.

DAUVOIS, S., S. LI, C. MARTEL and F. LABRIE (1989). "Inhibitory effect of androgens on DMBA-induced mammary carcinoma in the rat." Breast cancer research and treatment **14**(3): 299-306.

DAY, J., M., H. J. TUTILL, S. P. NEWMAN, A. PUROHIT, H. R. LAWRENCE, N. VICKER, B. V. POTTER and M. J. REED (2006). "17 β -Hydroxysteroid dehydrogenase Type 1 and Type 2: Association between mRNA expression and activity in cell lines." Molecular and cellular endocrinology **248**(1): 246-249.

DE BOSSCHER, K., K. VAN CRAENENBROECK, O. MEIJER, C. and G. HAEGEMAN (2008). "Selective transrepression versus transactivation mechanisms by glucocorticoid receptor modulators in stress and immune systems." European Journal of Pharmacology **583**(2–3): 290-302.

DILLER, M., S. SCHÜLER, S. BUCHHOLZ, C. LATTRICH, O. TREECK and O. ORTMANN (2014). "Effects of estriol on growth, gene expression and estrogen response element activation in human breast cancer cell lines." Maturitas **77**(4): 336-343.

DIMITRAKAKIS, C. (2011). "Androgens and Breast Cancer in Men and Women." Endocrinology and Metabolism Clinics of North America **40**(3): 533-547.

DIMITRAKAKIS, C. and C. BONDY (2009). "Androgens and the breast." Breast Cancer Research **11**(5): 212.

DIMITRAKAKIS, C., J. ZHOU, J. WANG, A. BELANGER, F. LABRIE, C. CHENG, D. POWELL and C. BONDY (2003). "A physiologic role for testosterone in limiting estrogenic stimulation of the breast." Menopause **10**(4): 292-298.

DOTZLAW, H., E. LEYGUE, P. H. WATSON and L. MURPHY, C. (1999). "Estrogen Receptor- β Messenger RNA Expression in Human Breast Tumor Biopsies: Relationship to Steroid Receptor Status and Regulation by Progestins." Cancer Research **59**(3): 529-532.

DRUMMOND, A., E. and P. FULLER, J. (2012). "Activin and inhibin, estrogens and NF κ B, play roles in ovarian tumourigenesis is there crosstalk?" Molecular and Cellular Endocrinology **359**(1-2): 85-91.

DUNN, J. F., B. C. NISULA and D. RODBARD (1981). "Transport of steroid hormones: binding of 21 endogenous steroids to both testosterone-binding globulin and corticosteroid-binding globulin in human plasma." The Journal of Clinical Endocrinology & Metabolism **53**(1): 58-68.

ENMARK, E. and J. Å. GUSTAFSSON (1999). "Oestrogen receptors – an overview." Journal of Internal Medicine **246**(2): 133-138.

ENMARK, E., P.-H. MARKKU, G. KAJ, L. SVETLANA, L. JACOB, F. GABRIEL, N. MAGNUS and G. JAN-ÅKE (1997). "Human Estrogen Receptor β -Gene Structure, Chromosomal Localization, and Expression Pattern." The Journal of Clinical Endocrinology & Metabolism **82**(12): 4258-4265.

FARMER, P., H. BONNEFOI, V. BECETTE, M. TUBIANA-HULIN, P. FUMOLEAU, D. LARSIMONT, G. MACGROGAN, J. BERGH, D. CAMERON, D. GOLDSTEIN, S. DUSS, A.-L. NICOULAZ, C. BRISKEN, M. FICHE, M. DELORENZI and R. IGGO (2005). "Identification of molecular apocrine breast tumours by microarray analysis." Oncogene **24**(29): 4660-4671.

FARQUHAR, C., J. MARJORIBANKS, A. LETHABY, J. A. SUCKLING and Q. LAMBERTS (2009). "Long term hormone therapy for perimenopausal and postmenopausal women." Cochrane Database of Systematic Reviews(2).

FAULDS, M., HEDENGRAN., C. ZHAO, K. DAHLMAN-WRIGHT and J.-Å. GUSTAFSSON (2012). "The diversity of sex steroid action: regulation of metabolism by estrogen signaling." Journal of Endocrinology **212**(1): 3-12.

FIELDS, S. and O.-K. SONG (1989). "A novel genetic system to detect protein protein interactions."

FIORETTI, F. M., A. SITA-LUMSDEN, C. L. BEVAN and G. N. BROOKE (2014). "Revising the role of the androgen receptor in breast cancer." Journal of Molecular Endocrinology **52**(3): R257-R265.

FORREST, D. and A. SWAROOP (2012). "Minireview: The Role of Nuclear Receptors in Photoreceptor Differentiation and Disease." Molecular Endocrinology **26**(6): 905-915.

FOWLER, A., M., N. SOLODIN, M. PREISLER-MASHEK, T., P. ZHANG, A. LEE, V. and E. ALARID, T. (2004). "Increases in estrogen receptor- α concentration in breast cancer cells promote serine 118/104/106-independent AF-1 transactivation and growth in the absence of estrogen." The FASEB Journal **18**(1): 81-93.

FOX, E., M., R. DAVIS, J. and M. SHUPNIK, A. (2008). "ER β in breast cancer—Onlooker, passive player, or active protector?" Steroids **73**(11): 1039-1051.

FRASOR, J., J. M. DANES, B. KOMM, K. C. N. CHANG, L. RICHARD, C. and B. S. KATZENELLENBOGEN (2003). "Profiling of estrogen up- and down-regulated gene expression in human breast cancer cells: Insights into gene networks and pathways underlying estrogenic control of proliferation and cell phenotype." Endocrinology **144**(10): 4562-4574.

FRASOR, J., A. WEAVER, M. PRADHAN, Y. DAI, L. D. MILLER, C. Y. LIN and A. STANCULESCU (2009). "Positive cross-talk between estrogen receptor and NF- κ B in breast cancer." Cancer Research **69**(23): 8918-8925.

FRESHNEY, R., I. (1987). "Animal Cell Culture: A Practical Approach. ." Journal of Basic Microbiology **27**(5): 274-274.

FULLER, P., J. (1991). "The steroid receptor superfamily: mechanisms of diversity." The FASEB Journal **5**(15): 3092-3099.

FURNESS, S., H. ROBERTS, J. MARJORIBANKS, A. LETHABY, M. HICKEY and C. FARQUHAR (2009). "Hormone therapy in postmenopausal women and risk of endometrial hyperplasia." Cochrane Database of Systematic Reviews(2).

GAGO, F., E., M. FANELLI, A. and D. CIOCCA, R. (2006). "Co-expression of steroid hormone receptors (estrogen receptor α and/or progesterone receptors) and Her2/neu (c-erbB-

2) in breast cancer: Clinical outcome following tamoxifen-based adjuvant therapy." The Journal of Steroid Biochemistry and Molecular Biology **98**(1): 36-40.

GALIEN, R. and T. GARCIA (1997). "Estrogen receptor impairs interleukin-6 expression by preventing protein binding on the NF- κ B site." Nucleic Acids Research **25**(12): 2424-2429.

GAO, W. and J. DALTON, T. (2007). "Expanding the therapeutic use of androgens via selective androgen receptor modulators (SARMs)." Drug Discovery Today **12**(5–6): 241-248.

GARAY, J., P. and B. PARK, H. (2012). "Androgen receptor as a targeted therapy for breast cancer." American Journal of Cancer Research **2**(4): 434-445.

GARCIA, M., D. DEROCQ, G. FREISS and H. ROCHEFORT (1992). "Activation of estrogen receptor transfected into a receptor-negative breast cancer cell line decreases the metastatic and invasive potential of the cells." Proceedings of the National Academy of Sciences of the United States of America **89**(23): 11538-11542.

GASC, J.-M. and E. E. BAULIEU (1987). "From the structure of steroid receptors to their assessment by immunocytochemistry in target cells." Journal of Steroid Biochemistry **27**(1–3): 177-184.

GELFAND, M. M. and A. FERENCZY (1989). "A prospective 1-year study of estrogen and progestin in postmenopausal women: Effects on the endometrium." Obstetrics and Gynecology **74**(3 PART 1): 398-402.

GHOSH, S., M. MAY, J. and E. KOPP, B. (1998). "NF- κ B and REL proteins: Evolutionarily Conserved Mediators of Immune Responses." Annual Review of Immunology **16**(1): 225-260.

GIACOBINI, E. (1998). "Aging, Alzheimer's disease, and estrogen therapy." Experimental Gerontology **33**(7–8): 865-869.

GIL, J., F., T. AUGUSTINE, N. and M. HOSIE, J. (2013). "Anastrozole and RU486: Effects on estrogen receptor α and Mucin 1 expression and correlation in the MCF-7 breast cancer cell line." Acta Histochemica **115**(8): 851-857.

GILLES, F., H. BRAND, S. DENGGER, R. METIVIER, M. KOS, G. REID, V. SONNTAG-BUCK and F. GANNON (2000). Identification of a new isoform of the human estrogen receptor alpha (hER- α) that is encoded by distinct transcripts and that is able to repress hER- α activation function 1. The EMBO Journal, The EMBO Journal.

GIRAULT, I., I. BIÈCHE and R. LIDEREAU (2006). "Role of estrogen receptor α transcriptional coregulators in tamoxifen resistance in breast cancer." Maturitas **54**(4): 342-351.

GIULIANELLI, S., A. MOLINOLO and C. LANARI (2013). Chapter Five - Targeting Progesterone Receptors in Breast Cancer. Vitamins & Hormones. L. Gerald, Academic Press. **Volume 93**: 161-184.

GLASER, R., L. and C. DIMITRAKAKIS (2013). "Reduced breast cancer incidence in women treated with subcutaneous testosterone, or testosterone with anastrozole: A prospective, observational study." Maturitas **76**(4): 342-349.

GOBINET, J., N. POUJOL and C. SULTAN (2002). "Molecular action of androgens." Molecular and Cellular Endocrinology **198**(1-2): 15-24.

GOUGELET, A., C. BOUCLIER, V. MARSAUD, S. MAILLARD, S. MUELLER, O., K. KORACH, S. and J.-M. RENOIR (2005). "Estrogen receptor α and β subtype expression and transactivation capacity are differentially affected by receptor-, hsp90- and immunophilin-ligands in human breast cancer cells." The Journal of Steroid Biochemistry and Molecular Biology **94**(1-3): 71-81.

GOUGELET, A., S. MUELLER, O., K. KORACH, S. and J.-M. RENOIR (2007). "Oestrogen receptors pathways to oestrogen responsive elements: The transactivation function-1 acts as the keystone of oestrogen receptor (ER) β -mediated transcriptional repression of ER α ." The Journal of Steroid Biochemistry and Molecular Biology **104**(3-5): 110-122.

GRAHAM, J., D., C. YEATES, R. BALLEINE, L., S. HARVEY, S., J. MILLIKEN, S., A. BILOUS, M. and C. CLARKE, L. (1996). "Progesterone receptor A and B protein expression in human breast cancer." Journal of Steroid Biochemistry and Molecular Biology **56**: 93-98.

GREEN, S., W. PHILIPPE, K. VIJAY, K. ANDREE, B. JEAN-MARC, A. PATRICK and C. PIERRE (1986). "Human oestrogen receptor cDNA: sequence, expression and homology to v-erb-A." Nature **320**: 134-139.

GREEVE, M., A., R. ALLAN, K., J. HARVEY, M. and J. BENTEL, M. (2004). "Inhibition of MCF-7 breast cancer cell proliferation by 5 α -dihydrotestosterone; a role for p21(Cip1/Waf1)." Journal of Molecular Endocrinology **32**(3): 793-810.

GRIEKSPoor, A., W. ZWART, J. NEEFJES and R. MICHALIDES (2007). "Visualizing the action of steroid hormone receptors in living cells." Nuclear Receptor Signaling **5**: e003.

GRODSTEIN, F., M. J. STAMPFER, J. E. MANSON, G. A. COLDITZ, W. C. WILLETT, B. ROSNER, F. E. SPEIZER and C. H. HENNEKENS (1996). "Postmenopausal estrogen and progestin use and the risk of cardiovascular disease." New England Journal of Medicine **335**(7): 453-461.

GROSSE, A., S. BARTSCH and A. BANIAHMAD (2012). "Androgen receptor-mediated gene repression." Molecular and cellular endocrinology **352**(1): 46-56.

GRUBER, C., J., W. TSCHUGGUEL, C. SCHNEEBERGER and J. HUBER, C. (2002). "Production and Actions of Estrogens." New England Journal of Medicine **346**(5): 340-352.

HACKENBERG, R., S. LÜTTCHENS, J. HOFMANN, R. KUNZMANN, F. HÖLZEL and K.-D. SCHULZ (1991). "Androgen Sensitivity of the New Human Breast Cancer Cell Line MFM-223." Cancer Research **51**(20): 5722-5727.

HAENDLER, B., I. SCHÜTTKE and W. SCHLEUNING, D. (2001). "Androgen receptor signalling: comparative analysis of androgen response elements and implication of heat-shock protein 90 and 14-3-3 η ." Molecular and Cellular Endocrinology **173**(1-2): 63-73.

HALL, J., M., J. COUSE, F. and K. KORACH, S. (2001). "The Multifaceted Mechanisms of Estradiol and Estrogen Receptor Signaling." Journal of Biological Chemistry **276**(40): 36869-36872.

HALL, J., M., D. MCDONNELL, P. and K. KORACH, S. (2002). "Allosteric Regulation of Estrogen Receptor Structure, Function, and Coactivator Recruitment by Different Estrogen Response Elements." Molecular Endocrinology **16**(3): 469-486.

HAMED, S. and S. MINDY, KURZER. (2015). "Estrogen metabolism and breast cancer." Cancer Letters **356**(2): 231-243.

HAN, G. (2014). "ER, PR and HER2 testing in breast cancer." Diagnostic Histopathology **20**(11): 440-445.

HAN, Y., L. YANG, F. SUAREZ-SAIZ, S. SAN-MARINA, J. CUI and M. MINDEN, D. (2008). "Wilms' Tumor 1 Suppressor Gene Mediates Antiestrogen Resistance via Down-Regulation of Estrogen Receptor- α Expression in Breast Cancer Cells." Molecular Cancer Research **6**(8): 1347-1355.

HARRINGTON, W., R., S. SHENG, D. BARNETT, H., L. PETZ, N., J. KATZENELLENBOGEN, A. and B. KATZENELLENBOGEN, S. (2003). "Activities of estrogen receptor alpha- and beta-selective ligands at diverse estrogen responsive gene sites mediating transactivation or transrepression." Molecular and Cellular Endocrinology **206**(1-2): 13-22.

HARTMAN, J., A. STROM and J.-Å. GUSTAFSSON (2012). "Current concepts and significance of estrogen receptor β in prostate cancer." Steroids **77**(12): 1262-1266.

HARTMAN, J., A. STRÖM and J.-Å. GUSTAFSSON (2009). "Estrogen receptor beta in breast cancer—Diagnostic and therapeutic implications." Steroids **74**(8): 635-641.

HELDRING, N., A. PIKE, S. ANDERSSON, J. MATTHEWS, G. CHENG, J. HARTMAN, M. TUJAGUE, A. STRÖM, E. TREUTER, M. WARNER and J.-Å. GUSTAFSSON (2007). Estrogen Receptors: How Do They Signal and What Are Their Targets.

HELGUERO, L., A., M. FAULDS, HEDENGRAN., J.-A. GUSTAFSSON and L.-A. HALDOSEN (2005). "Estrogen receptors alfa (ER[alpha]) and beta (ER[beta]) differentially regulate proliferation and apoptosis of the normal murine mammary epithelial cell line HC11." Oncogene **24**(44): 6605-6616.

HELGUERO, L. A., M. FAULDS, HEDENGRAN, J.-A. GUSTAFSSON and L.-A. HALDOSEN (2005). "Estrogen receptors alfa (ER[alpha]) and beta (ER[beta]) differentially regulate proliferation and apoptosis of the normal murine mammary epithelial cell line HC11." Oncogene **24**(44): 6605-6616.

HELSEN, C. and C. FRANK (2014). "Looking at nuclear receptors from a new angle." Molecular and Cellular Endocrinology **382**: 97-106.

HELSEN, C., S. KERKHOF, L. CLINCKEMALIE, L. SPANS, M. LAURENT, S. BOONEN, D. VANDERSCHUEREN and F. CLAESSENS (2012). "Structural basis for nuclear hormone receptor DNA binding." Molecular and Cellular Endocrinology **348**(2): 411-417.

HENDERSON, B., E. and H. FEIGELSON, SPENCER. (2000). "Hormonal carcinogenesis." Carcinogenesis **21**(3): 427-433.

HICKEY, T. E., J. L. L. ROBINSON, J. S. CARROLL and W. D. TILLEY (2012). "Minireview: The Androgen Receptor in Breast Tissues: Growth Inhibitor, Tumor Suppressor, Oncogene?" Molecular Endocrinology **26**(8): 1252-1267.

HODIS, H., N. and W. MACK, J. (2014). "Hormone replacement therapy and the association with coronary heart disease and overall mortality: Clinical application of the timing hypothesis." The Journal of Steroid Biochemistry and Molecular Biology **142**(0): 68-75.

HOLM, M., O. ANJA, K. NIELS and T. ANNE (2014). "Lifestyle influences on the association between pre-diagnostic hormone replacement therapy and breast cancer prognosis—Results from The Danish ‘Diet, Cancer and Health’ prospective cohort." Maturitas.

HOLMES, K. A., J. S. SONG, X. S. LIU, M. BROWN and J. S. CARROLL (2008). "Nkx3-1 and LEF-1 Function as Transcriptional Inhibitors of Estrogen Receptor Activity." Cancer Research **68**(18): 7380-7385.

HOPP, T., A., H. WEISS, L., S. HILSENBECK, G., Y. CUI, D. C. ALLRED, K. B. HORWITZ and S. FUQUA, A. W. (2004). "Breast Cancer Patients with Progesterone Receptor PR-A-Rich Tumors Have Poorer Disease-Free Survival Rates." Clinical Cancer Research **10**(8): 2751-2760.

HORWITZ, K., B., D. ZAVA, T., A. THILAGAR, K., E. JENSEN, M. and W. MCGUIRE, L. (1978). "Steroid Receptor Analyses of Nine Human Breast Cancer Cell Lines." Cancer Research **38**(8): 2434-2437.

HOWELL, A. (2006). "Fulvestrant ('Faslodex'): Current and future role in breast cancer management." Critical Reviews in Oncology/Hematology **57**(3): 265-273.

HOWELL, S. J., S. R. D. JOHNSTON and A. HOWELL (2004). "The use of selective estrogen receptor modulators and selective estrogen receptor down-regulators in breast cancer." Best Practice & Research Clinical Endocrinology & Metabolism **18**(1): 47-66.

HU, R., S. DAWOOD, M. D. HOLMES, L. C. COLLINS, S. J. SCHNITT, K. COLE, J. D. MAROTTI, S. E. HANKINSON, G. A. COLDITZ and R. M. TAMIMI (2011). "Androgen Receptor Expression and Breast Cancer Survival in Postmenopausal Women." Clinical cancer research : an official journal of the American Association for Cancer Research **17**(7): 1867-1874.

HUANG, B., M. WARNER and J.-Å. GUSTAFSSON (2014b). "Estrogen receptors in breast carcinogenesis and endocrine therapy." Molecular and Cellular Endocrinology(0).

HUANG, L., H. YAN, X. JIANG, Y. ZHANG, X. ZHANG, Y. JI, B. ZENG, B. XU, G. YIN, S. LEE, Y. YAN, X. MA and Y. PENG (2014). "Reference gene selection for quantitative real-time reverse-transcriptase PCR in orchardgrass subjected to various abiotic stresses." Gene **553**(2): 158-165.

HUANG, Z., F. GUENGERICH, P. and L. KAMINSKY, S. (1998). "16Alpha-hydroxylation of estrone by human cytochrome P4503A4/5." Carcinogenesis **19**(5): 867-872.

HUDSON, W., H., C. YOUN and E. ORTLUND, A. (2014). "Crystal Structure of the Mineralocorticoid Receptor DNA Binding Domain in Complex with DNA." PLoS ONE **9**(9): e107000.

HYDER, S., M., C. CHIAPPETTA and G. STANCEL, M. (1999). "Interaction of human estrogen receptors α and β with the same naturally occurring estrogen response elements." Biochemical Pharmacology **57**(6): 597-601.

INGEMAR, P. (2000). "Estrogens in the causation of breast, endometrial and ovarian cancers — evidence and hypotheses from epidemiological findings." The Journal of Steroid Biochemistry and Molecular Biology **74**(5): 357-364.

ISHIBASHI, H., S. TAKASHI, S. SATOSHI, M. TAKUYA, K. CHIKA, T. TOUICHIROU, S. MAKOTO, H. MASASHI, K. TAKASHI and S. HIRONOBU (2003). "Sex Steroid Hormone Receptors in Human Thymoma." The Journal of Clinical Endocrinology & Metabolism **88**(5): 2309-2317.

ISHIDA, M., T. MITSUI, M. IZAWA and J. ARITA (2010). "Absence of ligand-independent transcriptional activation of the estrogen receptor via the estrogen response element in pituitary lactotrophs in primary culture." The Journal of Steroid Biochemistry and Molecular Biology **118**(1–2): 93-101.

ISOLA, J. (1993). "Immunohistochemical demonstration of androgen receptor in breast cancer and its relationship to other prognostic factors." Journal of Pathology **170**(1): 31-35.

IWAO, K., Y. MIYOSHI, C. EGAWA, N. IKEDA and S. NOGUCHI (2000). "Quantitative analysis of estrogen receptor- β mRNA and its variants in human breast cancers." International Journal of Cancer **88**(5): 733-736.

JENSEN, E., V. and V. JORDAN, CRAIG. (2003). "The Estrogen Receptor: A Model for Molecular Medicine." Clinical Cancer Research **9**(6): 1980-1989.

JIANG, X., S. J. ELLISON, E. T. ALARID and D. J. SHAPIRO (2007). "Interplay between the levels of estrogen and estrogen receptor controls the level of the granzyme inhibitor, proteinase inhibitor 9 and susceptibility to immune surveillance by natural killer cells." Oncogene **26**: 4106-4114.

JIN, L. and Y. LI (2010). "Structural and functional insights into nuclear receptor signaling." Advanced Drug Delivery Reviews **62**(13): 1218-1226.

JORDAN, V. C. (2007). "SERMs: Meeting the Promise of Multifunctional Medicines." Journal of the National Cancer Institute **99**(5): 350-356.

JORDAN, V. C. and L. FORD, G. (2011). "Paradoxical Clinical Effect of Estrogen on Breast Cancer Risk: A "New" Biology of Estrogen-induced Apoptosis." Cancer Prevention Research **4**(5): 633-637.

JOZAN, S., C. MOURE, M. GILLOIS and F. BAYARD (1979). "Effects of estrone on cell proliferation of a human breast cancer (MCF-7) in long term tissue culture." Journal of Steroid Biochemistry **10**(3): 341-342.

KALAITZIDIS, D. and T. GILMORE, D. (2005). "Transcription factor cross-talk: the estrogen receptor and NF- κ B." Trends in Endocrinology & Metabolism **16**(2): 46-52.

KALKHOVEN, E., L. KWAKKENBOS-ISBRÜCKER, S. DE LAAT, W., P. VAN DER SAAG, T. and B. V. DER BURG (1994). "Synthetic progestins induce proliferation of breast tumor cell lines via the progesterone or estrogen receptor." Molecular and Cellular Endocrinology **102**(1–2): 45-52.

KALKHOVEN, E., S. WISSINK, P. VAN DER SAAG, T. and B. V. D. BURG (1996). "Negative Interaction between the RelA(p65) Subunit of NF- κ B and the Progesterone Receptor." Journal of Biological Chemistry **271**(11): 6217-6224.

KANDOUZ, M., A. LOMBET, J. PERROT, Y. , D. JACOB, S. CARVAJAL, K. A., W. ROSTENE, A. THERWATH and A. GOMPEL (1999). "Proapoptotic effects of antiestrogens, progestins and androgen in breast cancer cells." Journal of Steroid Biochemistry and Molecular Biology **69**(1-6): 463-471.

KARANTANOS, T., C. EVANS, P., B. TOMBAL, T. THOMPSON, C., R. MONTIRONI and W. ISAACS, B. (2015). "Understanding the Mechanisms of Androgen Deprivation Resistance in Prostate Cancer at the Molecular Level." European Urology **67**(3): 470-479.

KASSEL, O. and P. HERRLICH (2007). "Crosstalk between the glucocorticoid receptor and other transcription factors: Molecular aspects." Molecular and Cellular Endocrinology **275**(1-2): 13-29.

KASTNER, P., A. KRUST, B. TURCOTTE, U. STROPP, L. TORA, H. GRONEMEYER and P. CHAMBON (1990). "Two distinct estrogen-regulated promoters generate transcripts encoding the two functionally different human progesterone receptor forms A and B." The EMBO Journal **9**(5): 1603-1614.

KATALINIC, A. and R. RAWAL (2008). "Decline in breast cancer incidence after decrease in utilisation of hormone replacement therapy." Breast Cancer Research and Treatment **107**(3): 427-430.

KATZENELLENBOGEN, B., S. (1984). "Biology and receptor interactions of estriol and estriol derivatives in vitro and in vivo." Journal of Steroid Biochemistry **20**(4, Part 2): 1033-1037.

KATZENELLENBOGEN, B., S. and K. KORACH, S. (1997). "Editorial: A New Actor in the Estrogen Receptor Drama—Enter ER- β ." Endocrinology **138**(3): 861-862.

KHAN, J., A., C. BELLANCE, A. GUIOCHON-MANTEL, M. LOMBÈS and H. LOOSFELT (2012). "Differential Regulation of Breast Cancer-Associated Genes by Progesterone Receptor Isoforms PRA and PRB in a New Bi-Inducible Breast Cancer Cell Line." PLoS ONE **7**(9): e45993.

KIJIMA, I., T. ITOH and S. CHEN (2005). "Growth inhibition of estrogen receptor-positive and aromatase-positive human breast cancer cells in monolayer and spheroid cultures by letrozole, anastrozole, and tamoxifen." The Journal of Steroid Biochemistry and Molecular Biology **97**(4): 360-368.

- KIM, J. and G. A. COETZEE (2004). "Prostate specific antigen gene regulation by androgen receptor." Journal of cellular biochemistry **93**(2): 233-241.
- KITTLER, R., J. ZHOU, S. HUA, L. MA, Y. LIU, E. PENDLETON, C. CHENG, M. GERSTEIN and K. P. WHITE (2013). "A Comprehensive Nuclear Receptor Network for Breast Cancer Cells." Cell Reports **3**(2): 538-551.
- KLEIN-HITPASS, L., G. U. RYFFEL, E. HEITLINGER and A. CATO, C. B. (1988). "A 13 bp palindrome is a functional estrogen responsive element and interacts specifically with estrogen receptor." Nucleic Acids Research **16**(2): 647-663.
- KLOCK, G., U. STRAHLE and G. SCHUTZ (1987). "Oestrogen and glucocorticoid responsive elements are closely related but distinct." Nature **329**(6141): 734-736.
- KNOWER, K., C., S. TO, Q. and C. CLYNE, D. (2013). "Intracrine oestrogen production and action in breast cancer: An epigenetic focus." The Journal of Steroid Biochemistry and Molecular Biology **137**(0): 157-164.
- KNÜPFER, H. and R. PREIß (2007). "Significance of interleukin-6 (IL-6) in breast cancer (review)." Breast cancer research and treatment **102**(2): 129-135.
- KNUTSON, T., P. and C. LANGE, A. (2014). "Tracking progesterone receptor-mediated actions in breast cancer." Pharmacology & Therapeutics **142**(1): 114-125.
- KOCANOVA, S., E. A. KERR, S. RAFIQUE, S. BOYLE, E. KATZ, S. CAZE-SUBRA, W. A. BICKMORE and K. BYSTRICKY (2010). "Activation of Estrogen-Responsive Genes Does Not Require Their Nuclear Co-Localization." PLoS Genet **6**(4): e1000922.
- KOIDE, A., C. ZHAO, M. NAGANUMA, J. ABRAMS, S. DEIGHTON-COLLINS, D. SKAFAR, F. and S. KOIDE (2007). "Identification of Regions within the F Domain of the Human Estrogen Receptor α that Are Important for Modulating Transactivation and Protein-Protein Interactions." Molecular Endocrinology **21**(4): 829-842.
- KOTSOPOULOS, J. and S. NAROD, A. (2012). "Androgens and breast cancer." Steroids **77**(1-2): 1-9.
- KREBS, S., M. FISCHALECK and H. BLUM (2009). "A simple and loss-free method to remove TRIzol contaminations from minute RNA samples." Analytical Biochemistry **387**(1): 136-138.
- KUENEN-BOUMEESTER, V., T. H. VAN DER KWAST, C. C. CLAASSEN, M. P. LOOK, G. S. LIEM, J. G. M. KLIJN and S. C. HENZEN-LOGMANS (1996). "The clinical significance of androgen receptors in breast cancer and their relation to histological and cell biological parameters." European Journal of Cancer **32**(9): 1560-1565.

KUIPER, G., G., E. ENMARK, M. PELTO-HUIKKO, S. NILSSON and J. GUSTAFSSON, A. (1996). "Cloning of a novel receptor expressed in rat prostate and ovary." Proceedings of the National Academy of Sciences of the United States of America **93**(12): 5925-5930.

KUIPER, G. G. J. M., B. CARLSSON, K. GRANDIEN, E. ENMARK, J. HÄGGBLAD, S. NILSSON and J.-Å. K. GUSTAFSSON (1997). "Comparison of the Ligand Binding Specificity and Transcript Tissue Distribution of Estrogen Receptors ER α and β ." Endocrinology **138**(3): 863-870.

KULSHRESHTHA, P., A. CHAKRABORTY, L. SINGH, A. K. MISHRA, D. BHATNAGAR and S. SAXENA (2010). "Androgen receptor status predicts response to chemotherapy, not risk of breast cancer in Indian women." World journal of surgical oncology **8**(1): 64.

KUMAR, R. and G. LITWACK (2009). "Structural and functional relationships of the steroid hormone receptors' N-terminal transactivation domain." Steroids **74**(12): 877-883.

KUMAR, V. and P. CHAMBON (1988). "The estrogen receptor binds tightly to its responsive element as a ligand-induced homodimer." Cell **55**(1): 145-156.

LAKIS, S., V. KOTOULA, A. ELEFThERAKI, G., A. BATISTATOU, M. BOBOS, T. KOLETSA, E. TIMOTHEADOU, S. CHRISAFI, G. PENTHEROUDAKIS, A. KOUTRAS, F. ZAGOURI, H. LINARDOU and G. FOUNTZILAS (2014). "The androgen receptor as a surrogate marker for molecular apocrine breast cancer subtyping." The Breast **23**(3): 234-243.

LAMB, A., D., C. MASSIE, E. and D. NEAL, E. (2014). "The transcriptional programme of the androgen receptor (AR) in prostate cancer." BJU International **113**(3): 358-366.

LANARI, C., V. WARGON, P. ROJAS and A. MOLINOLO, A. (2012). "Antiprogesterins in breast cancer treatment: are we ready?" Endocrine-Related Cancer **19**(3): R35-R50.

LANGE, C., A. (2008). "Challenges to defining a role for progesterone in breast cancer." Steroids **73**(9-10): 914-921.

LANZINO, M., F. DE AMICIS, M. J. MCPHAUL, S. MARSICO, M. L. PANNO and S. ANDÒ (2005). "Endogenous Coactivator ARA70 Interacts with Estrogen Receptor α (ER α) and Modulates the Functional ER α /Androgen Receptor Interplay in MCF-7 Cells." Journal of Biological Chemistry **280**(21): 20421-20430.

LAZENNEC, G., D. BRESSON, A. LUCAS, C. CHAUVEAU and F. VIGNON (2001). "ER β Inhibits Proliferation and Invasion of Breast Cancer Cells." Endocrinology **142**(9): 4120-4130.

LAZENNEC, G. and B. KATZENELLENBOGEN, S. (1999). "Expression of human estrogen receptor using an efficient adenoviral gene delivery system is able to restore hormone-

dependent features to estrogen receptor-negative breast carcinoma cells." Molecular and Cellular Endocrinology **149**(1–2): 93-105.

LEE, C. H., Y.-T. JEON, S.-H. KIM and Y.-S. SONG (2007). "NF- κ B as a potential molecular target for cancer therapy." BioFactors **29**(1): 19-35.

LEE, H.-J. and C. CHANG (2003). "Recent advances in androgen receptor action." Cellular and Molecular Life Sciences CMLS **60**(8): 1613-1622.

LEE, H.-W., D.-H. AHN, S. CRAWLEY, C., J.-D. LI, J. GUM, R., C. BASBAUM, B., N. FAN, Q., D. SZYMKOWSKI, E., S.-Y. HAN, B. LEE, H., M. SLEISENGER, H. and Y. KIM, S. (2002). "Phorbol 12-Myristate 13-Acetate Up-regulates the Transcription of MUC2 Intestinal Mucin via Ras, ERK, and NF- κ B." Journal of Biological Chemistry **277**(36): 32624-32631.

LEE, W.-L., M.-H. CHENG, D.-C. TARNG, W.-C. YANG, F.-K. LEE and P.-H. WANG (2013). "The benefits of estrogen or selective estrogen receptor modulator on kidney and its related disease—chronic kidney disease—mineral and bone disorder: Osteoporosis." Journal of the Chinese Medical Association **76**(7): 365-371.

LEMON, H. M. (1975). "Estriol Prevention of Mammary Carcinoma Induced by 7,12-Dimethylbenzanthracene and Procarbazine." Cancer Research **35**(5): 1341-1353.

LEMON, H. M., P. F. KUMAR, C. PETERSON, J. F. RODRIGUEZ-SIERRA and K. M. ABBO (1989). "Inhibition of radiogenic mammary carcinoma in rats by estriol or tamoxifen." Cancer **63**(9): 1685-1692.

LEO, C., L. JOYCE., G. CHUNHUA, T. CHOW, WOON., E. SWEE, AW. and C. VALERIE, L, LIN. (2004). "Glucocorticoid and Mineralocorticoid Cross-Talk with Progesterone Receptor to Induce Focal Adhesion and Growth Inhibition in Breast Cancer Cells." Endocrinology **145**(3): 1314-1321.

LEUNG, B. S., A. H. POTTER and S. QURESHI (1981). "Interaction of prolactin, estrogen and progesterone in a human mammary carcinoma cell line, cama-1—I. cell growth and thymidine uptake." Journal of Steroid Biochemistry **15**: 421-427.

LEUNG, Y.-K., M.-T. LEE, H.-M. LAM, P. TARAPORE and S.-M. HO (2012). "Estrogen receptor-beta and breast cancer: Translating biology into clinical practice." Steroids **77**(7): 727-737.

LEVENSON, A., S. and V. C. JORDAN (1994). "Transfection of human estrogen receptor (ER) cDNA into ER-negative mammalian cell lines." The Journal of Steroid Biochemistry and Molecular Biology **51**(5–6): 229-239.

LEYGUE, E., H. DOTZLAW, P. H. WATSON and L. C. MURPHY (1998). "Altered Estrogen Receptor α and β Messenger RNA Expression during Human Breast Tumorigenesis." Cancer Research **58**(15): 3197-3201.

LI-MIN, S., L. JI-AN, C. SHIH-NI, S. FUNG-CHANG, M. CHIH-HSIN and K. CHIA-HUNG (2012). "Estrogen decrease coronary artery disease risk in patients with cervical cancer after treatment." gynecologic oncology **127**(1): 186-190.

LI, H., X. REN, E. LEBLANC, K. FREWIN, P. RENNIE, S. and A. CHERKASOV (2013). "Identification of Novel Androgen Receptor Antagonists Using Structure- and Ligand-Based Methods." Journal of Chemical Information and Modeling **53**(1): 123-130.

LI, Q. Q. and C. T. BEVER (2001). "Glatiramer acetate blocks interleukin-1-dependent nuclear factor- κ B activation and RANTES expression in human U-251 MG astroglial cells." Molecular Brain Research **87**(1): 48-60.

LI, R., J. CUI and Y. SHEN (2014). "Brain sex matters: Estrogen in cognition and Alzheimer's disease." Molecular and Cellular Endocrinology **389**(1-2): 13-21.

LIANG, Y., S.-Y. LIN, F. BRUNICARDI, CHARLES., J. GOSS and K. LI (2009). "DNA Damage Response Pathways in Tumor Suppression and Cancer Treatment." World Journal of Surgery **33**(4): 661-666.

LIBERMANN, T. A. and D. BALTIMORE (1990). "Activation of interleukin-6 gene expression through the NF-kappa B transcription factor." Molecular and Cellular Biology **10**(5): 2327-2334.

LIM, Y.-P., C.-L. LIN, Y.-N. LIN, W.-C. MA, D.-Z. HUNG and C.-H. KAO (2015). "Tamoxifen Treatment and the Reduced Risk of Hyperlipidemia in Asian Patients With Breast Cancer: A Population-Based Cohort Study." Clinical Breast Cancer(0).

LIN, C.-Y., S. ANDERS, S. L. KONG., S. KIETZ., S. JANE, THOMSEN., B. JASON, S. TEE., B. VINSENSIUS, VEGA., D. LANCE, MILLER., S. JOHANNA, B. JONAS, G. JAN-ÅKE and T. EDISON, LIU. (2007). "Inhibitory effects of estrogen receptor beta on specific hormone-responsive gene expression and association with disease outcome in primary breast cancer." Breast Cancer Research **9**(2).

LIN, H.-Y., M. SUN, C. LIN, H.-Y. TANG, D. LONDON, A. SHIH, F. DAVIS, B. and P. DAVIS, J. (2009). "Androgen-induced human breast cancer cell proliferation is mediated by discrete mechanisms in estrogen receptor- α -positive and -negative breast cancer cells." The Journal of Steroid Biochemistry and Molecular Biology **113**(3-5): 182-188.

LIN, H.-Y., M. SUN, C. LIN, H.-Y. TANG, D. LONDON, A. SHIH, F. B. DAVIS and P. J. DAVIS (2009). "Androgen-induced human breast cancer cell proliferation is mediated by

discrete mechanisms in estrogen receptor- \pm -positive and -negative breast cancer cells." The Journal of Steroid Biochemistry and Molecular Biology **113**(3&4): 182-188.

LIU, M.-M., C. ALBANESE, C. M. ANDERSON, K. HILTY, P. WEBB, R. M. UHT, R. H. PRICE, R. G. PESTELL and P. J. KUSHNER (2002). "Opposing action of estrogen receptors α and β on cyclin D1 gene expression." Journal of Biological Chemistry **277**(27): 24353-24360.

LOIBL, S., B. MÜLLER, G. VON MINCKWITZ, M. SCHWABE, M. ROLLER, S. DARB-ESFAHANI, B. ATASEVEN, A. DU BOIS, A. FISSLER-ECKHOFF, B. GERBER, U. KULMER, J.-U. ALLES, K. MEHTA and C. DENKERT (2011). "Androgen receptor expression in primary breast cancer and its predictive and prognostic value in patients treated with neoadjuvant chemotherapy." Breast Cancer Research and Treatment **130**(2): 477-487.

LONARD, D. M., Z. NAWAZ, C. L. SMITH and B. W. O'MALLEY (2000). "The 26S Proteasome Is Required for Estrogen Receptor- α and Coactivator Turnover and for Efficient Estrogen Receptor- α Transactivation." Molecular Cell **5**(6): 939-948.

LU, N. Z., S. E. WARDELL, K. L. BURNSTEIN, D. DEFRANCO, P. J. FULLER, V. GIGUERE, R. B. HOCHBERG, L. MCKAY, J.-M. RENOIR, N. L. WEIGEL, E. M. WILSON, D. P. MCDONNELL and J. A. CIDLOWSKI (2006). "International Union of Pharmacology. LXV. The Pharmacology and Classification of the Nuclear Receptor Superfamily: Glucocorticoid, Mineralocorticoid, Progesterone, and Androgen Receptors." Pharmacological Reviews **58**(4): 782-797.

LUBOSHITS, G., S. SHINA, O. KAPLAN, S. ENGELBERG, D. NASS, B. LIFSHITZ-MERCER, S. CHAITCHIK, I. KEYDAR and A. BEN-BARUCH (1999). "Elevated expression of the CC chemokine regulated on activation, normal T cell expressed and secreted (RANTES) in advanced breast carcinoma." Cancer research **59**(18): 4681-4687.

LYNN, J. (2004). "Fulvestrant ('Faslodex')—a new hormonal treatment for advanced breast cancer." European Journal of Oncology Nursing **8, Supplement 2**(0): S83-S88.

MACEDO, L. F., Z. GUO, S. L. TILGHMAN, G. J. SABNIS, Y. QIU and A. BRODIE (2006). "Role of Androgens on MCF-7 Breast Cancer Cell Growth and on the Inhibitory Effect of Letrozole." Cancer Research **66**(15): 7775-7782.

MACGREGOR, J. I. and V. C. JORDAN (1998). "Basic guide to the mechanisms of antiestrogen action." Pharmacological reviews **50**(2): 151-196.

MACLAUGHLIN, D. T. and G. S. RICHARDSON (1979). "Specificity of medroxyprogesterone acetate binding in human endometrium: Interaction with testosterone and progesterone binding sites." Journal of Steroid Biochemistry **10**(4): 371-377.

MAEDA, M. (2001). "The conserved residues of the ligand-binding domains of steroid receptors are located in the core of the molecules1." Journal of Molecular Graphics and Modelling **19**(6): 543-551.

MANGELSDORF, D., J., C. THUMMEL, M. BEATO, P. HERRLICH, G. SCHÜTZ, K. UMESONO, B. BLUMBERG, P. KASTNER, M. MARK, P. CHAMBON and R. EVANS, M. (1995). "The nuclear receptor superfamily: The second decade." Cell **83**(6): 835-839.

MARSDEN, J. (2003). "The menopause, hormone replacement therapy and breast cancer." Journal of Steroid Biochemistry and Molecular Biology **83**: 123-132.

MARTIN, L., A., S. PANCHOLI, C. CHAN, M. W., I. FARMER, C. KIMBERLEY, M. DOWSETT and S. JOHNSTON, R. D. (2005). "The anti-oestrogen ICI 182,780, but not tamoxifen, inhibits the growth of MCF-7 breast cancer cells refractory to long-term oestrogen deprivation through down-regulation of oestrogen receptor and IGF signalling." Endocrine-Related Cancer **12**(4): 1017-1036.

MATSUMOTO, T., M. SAKARI, M. OKADA, A. YOKOYAMA, S. TAKAHASHI, A. KOUZMENKO and S. KATO (2013). "The Androgen Receptor in Health and Disease." Annual Review of Physiology **75**(1): 201-224.

MATTER, C., M. and C. HANDSCHIN (2007). "RANTES (Regulated on Activation, Normal T Cell Expressed and Secreted), Inflammation, Obesity, and the Metabolic Syndrome." Circulation **115**(8): 946-948.

MATTHEWS, J. and J. A. GUSTAFSSON (2003). "Estrogen signaling: a subtle balance between ER alpha and ER beta." Mol Interv **3**(5): 281-292.

MCEWAN, I., J. (2012). "Nuclear hormone receptors: Allosteric switches." Molecular and Cellular Endocrinology **348**(2): 345-347.

MCKAY, L., I. and J. A. CIDLOWSKI. (2000). "CBP (CREB Binding Protein) Integrates NF- κ B (Nuclear Factor- κ B) and Glucocorticoid Receptor Physical Interactions and Antagonism." Molecular Endocrinology **14**(8): 1222-1234.

MCKAY, L., I. and J. CIDLOWSKI, A. (1999). "Molecular Control of Immune/Inflammatory Responses: Interactions Between Nuclear Factor- κ B and Steroid Receptor-Signaling Pathways." Endocrine Reviews **20**(4): 435-459.

MCKINLAY, S. M., D. J. BRAMBILLA and J. G. POSNER (1992). "The normal menopause transition." American journal of human biology **4**(1): 37-46.

MCNAMARA, K. M., N. L. MOORE, T. E. HICKEY, H. SASANO and W. D. TILLEY (2014b). "Complexities of androgen receptor signalling in breast cancer." Endocrine-Related Cancer **21**(4): T161-T181.

MCNAMARA, K. M., T. YODA, K. TAKAGI, Y. MIKI, T. SUZUKI and H. SASANO (2013). "Androgen receptor in triple negative breast cancer." The Journal of Steroid Biochemistry and Molecular Biology **133**(0): 66-76.

MCPHERSON, K., C. M. STEEL and J. M. DIXON (2000). "Breast cancer—epidemiology, risk factors, and genetics." BMJ : British Medical Journal **321**(7261): 624-628.

MEIYUN, F., M. B. ROBERT and P. N. KENNETH (2003). "The NEDD8 Pathway Is Required for Proteasome-Mediated Degradation of Human Estrogen Receptor (ER)- α and Essential for the Antiproliferative Activity of ICI 182,780 in ER α -Positive Breast Cancer Cells." Molecular Endocrinology **17**(3): 356-365.

MENASCE, L., G. WHITE, C. HARRISON and J. BOYLE (1993). "Localisation of the estrogen receptor locus (ESR) to chromosome 6q25.1 by FISH and a simple post-FISH banding technique." Genomics **17**: 263-265.

MIA, F. N., F. TOMMY, J. HEMMING, J. ULLA, R. LARS-ERIK, W. NILS and V. S. EVA (2013). "Hormone replacement therapy after breast cancer: 10 year follow up of the Stockholm randomised trial." European Journal of Cancer **49**: 52-59.

MILLER, W., R. and J. JACKSON (2003). "The therapeutic potential of aromatase inhibitors." Expert Opinion on Investigational Drugs **12**(3): 337-351.

MILLER, W. R. (2004). "Biological rationale for endocrine therapy in breast cancer." Best Practice & Research Clinical Endocrinology & Metabolism **18**(1): 1-32.

MIRZAI, S., L. CHUPANI, A. BARZEGARI, E., S. REZA, A. R. and M. JAMALAN (2013). "Novel inhibitor discovery against aromatase through virtual screening and molecular dynamic simulation: A computational approach in drug design." EXCLI Journal **12**: 168-183.

MISRAHI, M., H. LOOSFELT, M. ATGER, C. MERIEL, V. ZERAH, P. DESSEN and E. MILGROM (1988). "Organisation of the entire rabbit progesterone receptor mRNA and of the promoter and 5' flanking region of the gene." Nucleic acids research **16**(12): 5459-5472.

MITCHELL, S., H., P. MURTHA, E., S. ZHANG, W. ZHU and C. YOUNG, Y. F. (2000). "An androgen response element mediates LNCaP cell dependent androgen induction of the hK2 gene." Molecular and Cellular Endocrinology **168**(1-2): 89-99.

MOGGS, J. G., T. C. MURPHY, F. L. LIM, D. J. MOORE, R. STUCKEY, K. ANTROBUS, I. KIMBER and G. ORPHANIDES (2005). "Anti-proliferative effect of estrogen in breast

cancer cells that re-express ER α is mediated by aberrant regulation of cell cycle genes." Journal of Molecular Endocrinology **34**(2): 535-551.

MOINFAR, F., M. OKCU, O. TSYBROVSKYY, P. REGITNIG, S. F. LAX, W. WEYBORA, M. RATSCHKE, F. A. TAVASSOLI and H. DENK (2003). "Androgen receptors frequently are expressed in breast carcinomas." Cancer **98**(4): 703-711.

MONTANO, M. M., V. MULLER, A. TROBAUGH and B. S. KATZENELLENBOGEN (1995). "The carboxy-terminal F domain of the human estrogen receptor: role in the transcriptional activity of the receptor and the effectiveness of antiestrogens as estrogen antagonists." Molecular Endocrinology **9**(7): 814-825.

MORAS, D. and H. GRONEMEYER (1998). "The nuclear receptor ligand-binding domain: structure and function." Current Opinion in Cell Biology **10**(3): 384-391.

MORRISON, T. and J. WEISS (1998). Quantification of low-copy transcripts by continuous SYBR green I monitoring during amplification.

MOSSELMAN, S., J. POLMAN and R. DIJKEMA (1996). "ER β : Identification and characterization of a novel human estrogen receptor." FEBS Letters **392**(1): 49-53.

MRKLIĆ, I., Z. POGORELIĆ, V. ČAPKUN and S. TOMIĆ (2013). "Expression of androgen receptors in triple negative breast carcinomas." Acta Histochemica **115**(4): 344-348.

MULAC-JERICEVIC, B., J. LYDON, P., F. DEMAYO, J. and O. CONNEELY, M. (2003). "Defective mammary gland morphogenesis in mice lacking the progesterone receptor B isoform." Proceedings of the National Academy of Sciences of the United States of America **100**(17): 9744-9749.

MURPHY, E. (2011). "Estrogen Signaling and Cardiovascular Disease." Journal of American Heart Association **109****687-696**.

MURPHY, L., T. CHERLET, A. LEWIS, Y. BANU and P. WATSON (2003). "New insights into estrogen receptor function in human breast cancer." Annals of Medicine **35**(8): 614-631.

NANTASENAMAT, C., A. WORACHARTCHEEWAN, S. PRACHAYASITTIKUL, C. ISARANKURA-NA-AYUDHYA and V. PRACHAYASITTIKUL (2013). "QSAR modeling of aromatase inhibitory activity of 1-substituted 1,2,3-triazole analogs of letrozole." European Journal of Medicinal Chemistry **69**: 99-114.

NAVARRO, E., G. SERRANO-HERAS, M. J. CASTAÑO and J. SOLERA (2015). "Real-time PCR detection chemistry." Clinica Chimica Acta **439**(0): 231-250.

NEED, E. F., L. A. SELTH, T. J. HARRIS, S. N. BIRRELL, W. D. TILLEY and G. BUCHANAN (2012). "Research resource: Interplay between the genomic and transcriptional networks of androgen receptor and estrogen receptor α in luminal breast cancer cells." Molecular Endocrinology **26**(11): 1941-1952.

NELSON, E., R., S. WARDELL, E. and D. MCDONNELL, P. (2013). "The molecular mechanisms underlying the pharmacological actions of estrogens, SERMs and oxysterols: Implications for the treatment and prevention of osteoporosis." Bone **53**(1): 42-50.

NELSON, P. S., N. CLEGG, H. ARNOLD, C. FERGUSON, M. BONHAM, J. WHITE, L. HOOD and B. LIN (2002). "The program of androgen-responsive genes in neoplastic prostate epithelium." Proceedings of the National Academy of Sciences **99**(18): 11890-11895.

NEWMAN, S., P., C. IRESON, R., H. TUTILL, J., J. DAY, M., M. F. PARSONS, M. P. LEESE, B. V. POTTER, M. J. REED and A. PUROHIT (2006). "The role of 17 β -hydroxysteroid dehydrogenases in modulating the activity of 2-methoxyestradiol in breast cancer cells." Cancer research **66**(1): 324-330.

NEWTON, R. and N. HOLDEN, S. (2007). "Separating Transrepression and Transactivation: A Distressing Divorce for the Glucocorticoid Receptor?" Molecular Pharmacology **72**(4): 799-809.

NICHOLAS, S., AGOFF., E. PAUL, SWANSON., L. HANNAH, E. STEPHEN, HAWES. and J. THOMAS, LAWTON. (2003). "Androgen Receptor Expression in Estrogen Receptor-Negative Breast Cancer Immunohistochemical, Clinical, and Prognostic Associations." American Journal of Clinical Pathology **120**: 725-731.

NICOLÁS, D.-C., B., R. GERMÁN, F., A. GONZÁLEZ, R. RAMÍREZ, C. BILBAO, A. CABRERA DE LEÓN, J. AGUIRRE, A., R. CHIRINO, D. NAVARRO and J. DÍAZ-CHICO, C. (2007). "Androgens and androgen receptors in breast cancer." The Journal of Steroid Biochemistry and Molecular Biology **105**(1-5): 1-15.

NICOLÁS DÍAZ-CHICO, B., F. GERMÁN RODRÍGUEZ, A. GONZÁLEZ, R. RAMÍREZ, C. BILBAO, A. CABRERA DE LEÓN, A. AGUIRRE JAIME, R. CHIRINO, D. NAVARRO and J. C. DÍAZ-CHICO (2007). "Androgens and androgen receptors in breast cancer." The Journal of Steroid Biochemistry and Molecular Biology **105**(1-5): 1-15.

NILSEN, J., G. MOR and F. NAFTOLIN (2000). "Estrogen-regulated developmental neuronal apoptosis is determined by estrogen receptor subtype and the Fas/Fas ligand system." Journal of neurobiology **43**(1): 64-78.

NILSSON, S., S. MÄKELÄ, E. TREUTER, M. TUJAGUE, J. THOMSEN, G. ANDERSSON, E. ENMARK, K. PETTERSSON, M. WARNER and J.-Å. GUSTAFSSON (2001). Mechanisms of Estrogen Action.

NIU, J., Y. SHI, G. TAN, C. H. YANG, M. FAN, L. M. PFEFFER and Z.-H. WU (2012). "DNA Damage Induces NF- κ B-dependent MicroRNA-21 Up-regulation and Promotes Breast Cancer Cell Invasion." Journal of Biological Chemistry **287**(26): 21783-21795.

O'NEILL, P. A., M. P. A. DAVIES, A. M. SHAABAN, H. INNES, A. TOREVELL, D. R. SIBSON and C. S. FOSTER (2004). "Wild-type oestrogen receptor beta (ER[beta]1) mRNA and protein expression in Tamoxifen-treated post-menopausal breast cancers." Br J Cancer **91**(9): 1694-1702.

O'REGAN, R. M. and V. C. JORDAN (2002). "The evolution of tamoxifen therapy in breast cancer: selective oestrogen-receptor modulators and downregulators." The Lancet Oncology **3**(4): 207-214.

OBR, A. E. and D. P. EDWARDS (2012). "The biology of progesterone receptor in the normal mammary gland and in breast cancer." Molecular and Cellular Endocrinology **357**(1-2): 4-17.

OGAWA, S., S. INOUE, T. WATANABE, H. HIROI, A. ORIMO, T. HOSOI, Y. OUCHI and M. MURAMATSU (1998). "The Complete Primary Structure of Human Estrogen Receptor β (hER β) and Its Heterodimerization with ER α in Vivo and in Vitro " Biochemical and Biophysical Research Communications **243**(1): 122-126.

OMOTO, Y., S. INOUE, S. OGAWA, T. TOYAMA, H. YAMASHITA, M. MURAMATSU, S. KOBAYASHI and H. IWASE (2001). "Clinical value of the wild-type estrogen receptor β expression in breast cancer." Cancer Letters **163**(2): 207-212.

OSBORNE, C., KENT., R. SCHIFF, S. FUQUA, A. W. and J. SHOU (2001). "Estrogen Receptor: Current Understanding of Its Activation and Modulation." Cancer Research **7**(12 Supplement): 4338s-4342s.

OSBORNE, C. K., K. HOBBS and G. M. CLARK (1985). "Effect of Estrogens and Antiestrogens on Growth of Human Breast Cancer Cells in Athymic Nude Mice." Cancer Research **45**(2): 584-590.

OZGYIN, L., E. ERDÖS, D. BOJCSUK and B. BALINT, L. (2015). "Nuclear receptors in transgenerational epigenetic inheritance." Progress in Biophysics and Molecular Biology(0).

PAECH, K., P. WEBB, G. G. J. M. KUIPER, S. NILSSON, J.-Å. GUSTAFSSON, P. J. KUSHNER and T. S. SCANLAN (1997). "Differential Ligand Activation of Estrogen Receptors ER α and ER β at AP1 Sites." Science **277**(5331): 1508-1510.

PALVIMO, J., J., P. REINIKAINEN, T. IKONEN, P. KALLIO, J., A. MOILANEN and O. JÄNNE, A. (1996). "Mutual Transcriptional Interference between RelA and Androgen Receptor." Journal of Biological Chemistry **271**(39): 24151-24156.

PANET-RAYMOND, V., B. GOTTLIEB, L. BEITEL, K., L. PINSKY and M. TRIFIRO, A. (2000). "Interactions between androgen and estrogen receptors and the effects on their transactivational properties." Molecular and Cellular Endocrinology **167**(1&2): 139-150.

PANG, H. and E. FABER, LEE. (2001). "Estrogen and rapamycin effects on cell cycle progression in T47D breast cancer cells." Breast cancer research **70**: 21-26.

PARANJAPPE, S. G., A. V. TURANKAR, S. L. WAKODE and G. N. DAKHALE (2005). "Estrogen protection against coronary heart disease: Are the relevant effects of estrogen mediated through its effects on uterus – such as the induction of menstruation, increased bleeding, and the facilitation of pregnancy?" Medical Hypotheses **65**(4): 725-727.

PARDRIDGE, W., M. (1988). "Selective Delivery of Sex Steroid Hormones to Tissues In Vivo by Albumin and by Sex Hormone-Binding Globulina." Annals of the New York Academy of Sciences **538**(1): 173-192.

PARIS, O., L. FERRARO, O. M. V. GROBER, M. RAVO, M. R. DE FILIPPO, G. GIURATO, G. NASSA, R. TARALLO, C. CANTARELLA, F. RIZZO, A. DI BENEDETTO, M. MOTTOLESE, V. BENES, C. AMBROSINO, E. NOLA and A. WEISZ (2012). "Direct regulation of microRNA biogenesis and expression by estrogen receptor beta in hormone-responsive breast cancer." Oncogene **31**(38): 4196-4206.

PARK, S., J. KOO, H. S. PARK, J. H. KIM, S. Y. CHOI, J. H. LEE, B. W. PARK and K. S. LEE (2010). "Expression of androgen receptors in primary breast cancer." Annals of Oncology **21**(3): 488-492.

PARK, S., J. S. KOO, M. S. KIM, H. S. PARK, J. S. LEE, S. I. KIM, B. W. PARK and K. S. LEE (2011). "Androgen receptor expression is significantly associated with better outcomes in estrogen receptor-positive breast cancers." Annals of oncology: 678.

PARK, S., H. PARK, J. KOO, W. YANG, S. KIM and B.-W. PARK (2012). "Higher expression of androgen receptor is a significant predictor for better endocrine-responsiveness in estrogen receptor-positive breast cancers." Breast Cancer Research and Treatment **133**(1): 311-320.

PARUTHIYIL, S., H. PARMAR, V. KEREKATTE, G. CUNHA, R., G. FIRESTONE, L. and D. LEITMAN, C. (2004). "Estrogen Receptor ER β Inhibits Human Breast Cancer Cell Proliferation and Tumor Formation by Causing a G2 Cell Cycle Arrest." Cancer Research **64**(1): 423-428.

PASCUAL, G. and C. GLASS, K. (2006). "Nuclear receptors versus inflammation: mechanisms of transrepression." Trends in Endocrinology & Metabolism **17**(8): 321-327.

PASQUALINI, J. R. (2003). "Differential effects of progestins on breast tissue enzymes." Maturitas **46, Supplement 1**: 45-54.

PATANI, N. and L. MARTIN, A. (2014). "Understanding response and resistance to oestrogen deprivation in ER-positive breast cancer." Molecular and Cellular Endocrinology **382**(1): 683-694.

PEARCE, S., TIMM. and V. JORDAN, CRAIG. (2004). "The biological role of estrogen receptors α and β in cancer." Critical Reviews in Oncology/Hematology **50**(1): 3-22.

PERISSI, V. and M. ROSENFELD, G. (2005). "Controlling nuclear receptors: the circular logic of cofactor cycles." Nat Rev Mol Cell Biol **6**(7): 542-554.

PETERS, A. A., G. BUCHANAN, C. RICCIARDELLI, T. BIANCO-MIOTTO, M. M. CENTENERA, J. M. HARRIS, S. JINDAL, D. SEGARA, L. JIA, N. L. MOORE, S. M. HENSHALL, S. N. BIRRELL, G. A. COETZEE, R. L. SUTHERLAND, L. M. BUTLER and W. D. TILLEY (2009). "Androgen receptor inhibits estrogen receptor- α activity and is prognostic in breast cancer." Cancer Research **69**(15): 6131-6140.

PETRA, P. H. (1991). "The plasma sex steroid binding protein (SBP or SHBG). A critical review of recent developments on the structure, molecular biology and function." The Journal of Steroid Biochemistry and Molecular Biology **40**(4-6).

PETZ, L. N., Y. S. ZIEGLER, J. R. SCHULTZ, H. KIM, J. K. KEMPER and A. M. NARDULLI (2004). "Differential regulation of the human progesterone receptor gene through an estrogen response element half site and Sp1 sites." The Journal of steroid biochemistry and molecular biology **88**(2): 113-122.

PFAFFL, M., W. (2001). "A new mathematical model for relative quantification in real-time RT-PCR." Nucleic Acids Research **29**(9): e45.

PHILLIPS, M., A., J. D'AURIA, C., K. LUCK and J. GERSHENZON (2009). "Evaluation of candidate reference genes for real-time quantitative PCR of plant samples using purified cDNA as template." Plant Molecular Biology Reporter **27**(3): 407-416.

PIKE, M., C., R. PETERS, K., W. COZEN, N. PROBST-HENSCH, M., P. WAN, C., T. MACK, M. and J. FELIX, C. (1997). "Estrogen-Progestin Replacement Therapy and Endometrial Cancer." Journal of the National Cancer Institute **89**(15): 1110-1116.

PINTO, A., B., E. BINDER, F., W. KOHRT, M., D. BRONDER, R. and D. WILLIAMS, B. (2003). "Effects of trimonthly progestin administration on the endometrium in elderly postmenopausal women who receive hormone replacement therapy: A pilot study." American Journal of Obstetrics and Gynecology **189**(1): 11-15.

PISTELLI, M., M. CARAMANTI, T. BISCOTTI, A. SANTINELLI, A. PAGLIACCI, M. DE LISA, Z. BALLATORE, F. RIDOLFI, E. MACCARONI, R. BRACCI, R. BERARDI, N.

BATTELLI and S. CASCINU (2014). "Androgen receptor expression in early triple-negative breast cancer: Clinical significance and prognostic associations." Cancers **6**(3): 1351-1362.

PLAISANCE, S., V. B. WIM, B. ELKE, F. WALTER and H. GUY (1997). "Recombination Signal Sequence Binding Protein Jk Is Constitutively Bound to the NF- κ B Site of the Interleukin-6 Promoter and Acts as a Negative Regulatory Factor." Molecular and Cellular Biology **17**(7): 3733-3743.

POWELL, E., S. ERIN, B. ASHLEY, L. JUN, K. SUNDUZ, B. KARI, WISINSKI., H. WEI and X. WEI (2012). "Identification of estrogen receptor dimer selective ligands reveals growth-inhibitory effects on cells that co-express ER α and ER β ." PLoS One **7**(2).

PRATT, W., B. and D. TOFT, O. (2003). "Regulation of Signaling Protein Function and Trafficking by the hsp90/hsp70-Based Chaperone Machinery." Experimental Biology and Medicine **228**(2): 111-133.

QU, Q., Y. MAO, X.-C. FEI and K.-W. SHEN (2013). "The Impact of Androgen Receptor Expression on Breast Cancer Survival: A Retrospective Study and Meta-Analysis." PLoS ONE **8**(12): e82650.

RAVI, R. and A. BEDI (2004). "NF- κ B in cancer—a friend turned foe." Drug Resistance Updates **7**(1): 53-67.

RAVN, S., H. , J. ROSENBERG and E. BOSTOFTE (1994). "Postmenopausal hormone replacement therapy - clinical implications." European Journal of Obstetrics and Gynecology and Reproductive Biology **53**(2): 81-93.

RAY, P., S. GHOSH, K., D.-H. ZHANG and A. RAY (1997). "Repression of interleukin-6 gene expression by 17 β -estradiol: inhibition of the DNA-binding activity of the transcription factors NF-IL6 and NF- κ B by the estrogen receptor." FEBS Letters **409**(1): 79-85.

RECHOUM, Y., D. ROVITO, D. IACOPETTA, I. BARONE, S. ANDÒ, N. WEIGEL, B. O'MALLEY, P. BROWN and S. W. FUQUA (2014). "AR collaborates with ER α in aromatase inhibitor-resistant breast cancer." Breast Cancer Research and Treatment **147**(3): 473-485.

RENOIR, J.-M. (2012). "Estradiol receptors in breast cancer cells: Associated co-factors as targets for new therapeutic approaches." Steroids **77**(12): 1249-1261.

RIBEIRO, J., R. and R. FREIMAN, N. (2014). "Estrogen signaling crosstalk: Implications for endocrine resistance in ovarian cancer." The Journal of Steroid Biochemistry and Molecular Biology **143**(0): 160-173.

RICKARD, D. J., K. M. WATERS, T. J. RUESINK, S. KHOSLA, J. A. KATZENELLENBOGEN, B. S. KATZENELLENBOGEN, B. L. RIGGS and T. C.

SPELSBERG (2002). "Estrogen Receptor Isoform-Specific Induction of Progesterone Receptors in Human Osteoblasts." Journal of Bone and Mineral Research **17**(4): 580-592.

RIRIE, K., M., R. RASMUSSEN, P. and C. WITTEWER, T. (1997). "Product Differentiation by Analysis of DNA Melting Curves during the Polymerase Chain Reaction." Analytical Biochemistry **245**(2): 154-160.

RIZZA, P., B. INES, Z. DOMENICO, G. FRANCESCA, L. MARILENA, D. A. FRANCESCA, M. LOREDANA, S. DIEGO, C. STEFANIA, D. KARIN, WRIGHT., G. JAN-AKE and A. SEBASTIANO (2014). "Estrogen receptor beta as a novel target of androgen receptor action in breast cancer cell lines." Breast Cancer Research **16**.

ROBERTSON, S., F. ALLIE-REID, W. V. BERGHE, K. VISSER, A. BINDER, D. AFRICANDER, M. VISMER, K. DE BOSSCHER, J. HAPGOOD, G. HAEGEMAN and A. LOUW (2010). "Abrogation of Glucocorticoid Receptor Dimerization Correlates with Dissociated Glucocorticoid Behavior of Compound A." Journal of Biological Chemistry **285**(11): 8061-8075.

ROBINSON, J. L., S. MACARTHUR, C. S. ROSS-INNES, W. D. TILLEY, D. E. NEAL, I. G. MILLS and J. S. CARROLL (2011). "Androgen receptor driven transcription in molecular apocrine breast cancer is mediated by FoxA1." The EMBO journal **30**(15): 3019-3027.

ROGER, P., M. E. SAHLA, S. MÄKELÄ, J. Å. GUSTAFSSON, P. BALDET and H. ROCHEFORT (2001). "Decreased expression of estrogen receptor β protein in proliferative preinvasive mammary tumors." Cancer research **61**(6): 2537-2541.

ROSSOUW, E. J., G. L. ANDERSON., R. L. PRENTICE., A. Z. LACROIX., C. KOOPERBERG., M. L. STEFANICK., R. D. JACKSON., S. A. A. BERESFORD., B. V. HOWARD., K. C. JOHNSON., J. M. KOTCHEN. and J. OCKENE. (2002). "Risks and benefits of estrogen plus progestin in healthy postmenopausal women: Principal results from the women's health initiative randomized controlled trial." JAMA **288**(3): 321-333.

RUSSO, J., X. AO, C. GRILL and I. H. RUSSO (1999). "Pattern of distribution of cells positive for estrogen receptor α and progesterone receptor in relation to proliferating cells in the mammary gland." Breast Cancer Research and Treatment **53**(3): 217-227.

SABATIER, R., A. GONÇALVES and F. BERTUCCI (2014). "Personalized medicine: Present and future of breast cancer management." Critical Reviews in Oncology/Hematology **91**(3): 223-233.

SADAR, M. D., M. HUSSAIN and N. BRUCHOVSKY (1999). "Prostate cancer: molecular biology of early progression to androgen independence." Endocrine-related cancer **6**(4): 487-502.

SAITOH, M., O. MASAHIDE, T. KAZUHIRO, K. JUN, O. TSUYOSHI, D. MASAKAZU, T. TOSHIFUMI, I. I. HIDEK, M.-A. AKIKO, D. BOTAO, T. SEIJI and K. HIROHISA (2005). "Medroxyprogesterone Acetate Induces Cell Proliferation through Up-Regulation of Cyclin D1 Expression via Phosphatidylinositol 3-Kinase/Akt/Nuclear Factor-kB Cascade in Human Breast Cancer Cells." Endocrinology **146**(11): 4917-4925.

SAJI, S., H. MAKIKO and T. MASAKAZU (2005). "Clinical significance of estrogen receptor β in breast cancer." Cancer Chemother Pharmacol **56**(1): 21-26.

SAMAVAT, H. and M. S. KURZER (2015). "Estrogen Metabolism and Breast Cancer." Cancer Letters **356**(2 Part A): 231-243.

SAMBROOK, J., T. MANIATUS and E. FRITSCH, F. (1989). Molecular cloning: A Laboratory Manual(New York): Cold Spring Harbor Laboratory Press.

SANTEN, R., E. CAVALIERI, E. ROGAN, J. RUSSO, J. GUTTENPLAN, J. INGLE and W. YUE (2009). "Estrogen Mediation of Breast Tumor Formation Involves Estrogen Receptor-Dependent, as Well as Independent, Genotoxic Effects." Annals of the New York Academy of Sciences **1155**(1): 132-140.

SAXENA, R., I. FATIMA, V. CHANDRA, C. BLESSON, S., G. KHARKWAL, M. HUSSAIN, K., K. HAJELA, B. ROY, G. and A. DWIVEDI (2013). "Benzopyran derivative CDRI-85/287 induces G2-M arrest in estrogen receptor-positive breast cancer cells via modulation of estrogen receptors α - and β -mediated signaling, in parallel to EGFR signaling and suppresses the growth of tumor xenograft." Steroids **78**(11): 1071-1086.

SCARPATI, E., M. and J. SADLER, E. (1989). "Regulation of endothelial cell coagulant properties. Modulation of tissue factor, plasminogen activator inhibitors, and thrombomodulin by phorbol 12-myristate 13-acetate and tumor necrosis factor." Journal of Biological Chemistry **264**(34): 20705-20713.

SCHIPPINGER, W., P. REGITNIG, N. DANDACHI, K.-D. WERNECKE, T. BAUERNHOFER, H. SAMONIGG and F. MOINFAR (2006). "Evaluation of the prognostic significance of androgen receptor expression in metastatic breast cancer." Virchows Archiv **449**(1): 24-30.

SCHULTZ, J. R., L. N. PETZ and A. M. NARDULLI (2003). "Estrogen receptor α and Sp1 regulate progesterone receptor gene expression." Molecular and Cellular Endocrinology **201**(1-2): 165-175.

SCHULTZ, J. R., L. N. PETZ and A. M. NARDULLI (2005). "Cell- and Ligand-specific Regulation of Promoters Containing Activator Protein-1 and Sp1 Sites by Estrogen Receptors α and β ." Journal of Biological Chemistry **280**(1): 347-354.

SEOK-JUN, K., J.-Y. SHIN., L. KANG-DUCK, B. YOUNG-KI, S. KI WOONG, J. N. SEOK and C. KYUNG-HEE (2012). "MicroRNA let-7a suppresses breast cancer cell migration and invasion through downregulation of C-C chemokine receptor type 7." Breast Cancer Research **14**(1): R14.

SHAABAN, A. M., P. A. O'NEILL, M. P. A. DAVIES, R. SIBSON, C. R. WEST, P. H. SMITH and C. S. FOSTER (2003). "Declining estrogen receptor- β expression defines malignant progression of human breast neoplasia." The American journal of surgical pathology **27**(12): 1502-1512.

SHAFFER, P. L. and D. T. GEWIRTH (2002). "Structural basis of VDR-DNA interactions on direct repeat response elements." EMBO Journal **21**(9): 2242-2252.

SHAFIE, S. M. and L. A. LIOTTA (1980). "Formation of metastasis by human breast carcinoma cells (MCF-7) in nude mice." Cancer Letters **11**(2): 81-87.

SHAH, P., D., A. GUCALP and T. TRAINA, A. (2013). "The role of the androgen receptor in triple-negative breast cancer." Women's Health **9**(4): 351-360.

SIITERI, P., K. (1987). "Adipose tissue as a source of hormones." The American Journal of Clinical Nutrition **45**(1): 277-282.

SILVA, E., A. KABIL and A. KORTENKAMP (2010). "Cross-talk between non-genomic and genomic signalling pathways — Distinct effect profiles of environmental estrogens." Toxicology and Applied Pharmacology **245**(2): 160-170.

SILVERMANA, G., BARBARA., S.-D. NAVA, B. RONY and S. EHUD, KOKIAA. (2011). "Trends in breast cancer incidence associated with reductions in the use of hormone replacement therapy." Cancer Epidemiology Biomarkers & Prevention **35**: 11-16.

SIMONCINI, T., P. MANNELLA, L. FORNARI, A. CARUSO, G. VARONE and A. R. GENAZZANI (2004). "Genomic and non-genomic effects of estrogens on endothelial cells." Steroids **69**(8-9): 537-542.

SIMPSON, E., R. (2002). "Aromatization of androgens in women: current concepts and findings." Fertility and Sterility **77**, Supplement 4(0): 6-10.

SIVIK, T. and A. JANSSON (2012). "Progesterone and levonorgestrel regulate expression of 17 β HSD-enzymes in progesterone receptor positive breast cancer cell line T47D." Biochemical and Biophysical Research Communications **422**(1): 109-113.

SKLIRIS, G. P., E. LEYGUE, L. CURTIS-SNELL, P. H. WATSON and L. C. MURPHY (2006). "Expression of oestrogen receptor-[beta] in oestrogen receptor-[alpha] negative human breast tumours." British Journal of Cancer **95**(5): 616-626.

SKLIRIS, G. P., E. LEYGUE, P. H. WATSON and L. C. MURPHY (2008). "Estrogen receptor alpha negative breast cancer patients: Estrogen receptor beta as a therapeutic target." The Journal of Steroid Biochemistry and Molecular Biology **109**(1–2): 1-10.

SKOR, M. N., E. L. WONDER, M. KOCHERGINSKY, A. GOYAL, B. A. HALL, Y. CAI and S. D. CONZEN (2013). "Glucocorticoid receptor antagonism as a novel therapy for triple-negative breast cancer." Clinical Cancer Research **19**(22): 6163-6172.

SOMBOONPORN, W. and S. DAVIS, R. (2004). "Testosterone Effects on the Breast: Implications for Testosterone Therapy for Women." Endocrine Reviews **25**(3): 374-388.

SOMMER, S. and S. FUQUA, A. W. (2001). "Estrogen receptor and breast cancer." Seminars in Cancer Biology **11**(5): 339-352.

SØNDER, S., ULRIK, SALLING., A. WOETMANN, N. ØDUM and K. BENDTZEN (2006). "Spironolactone induces apoptosis and inhibits NF- κ B independent of the mineralocorticoid receptor." Apoptosis **11**(12): 2159-2165.

SONG, R. X.-D. and R. J. SANTEN (2006). "Membrane Initiated Estrogen Signaling in Breast Cancer." Biology of Reproduction **75**(1): 9-16.

SONG, R. X., Z. ZHANG and R. J. SANTEN (2005). "Estrogen rapid action via protein complex formation involving ER α and Src." Trends in Endocrinology & Metabolism **16**(8): 347-353.

SONG, R. X. D. and R. J. SANTEN (2003). "Apoptotic action of estrogen." Apoptosis **8**(1): 55-60.

SØREIDE, J., A., O. LEA, A., J. VARHAUG, E., A. SKARSTEIN and S. KVINNSLAND (1992). "Androgen receptors in operable breast cancer: relation to other steroid hormone receptors, correlations to prognostic factors and predictive value for effect of adjuvant tamoxifen treatment." European Journal Surgical Oncology. **18**(2): 112-118.

SOURLA, A., C. MARTEL, C. LABRIE and F. LABRIE (1998). "Almost exclusive androgenic action of dehydroepiandrosterone in the rat mammary gland." Endocrinology **139**(2): 753-764.

SPEIRS, V., P. CARDER, J., S. LANE, D. DODWELL, M. LANSDOWN, R. J. and A. HANBY, M. (2004). "Oestrogen receptor β : what it means for patients with breast cancer." The Lancet Oncology **5**(3): 174-181.

SPEIRS, V., C. MALONE, D. WALTON, S., M. KERIN, J. and S. ATKIN, L. (1999). "Increased Expression of Estrogen Receptor β mRNA in Tamoxifen-resistant Breast Cancer Patients." Cancer Research **59**(21): 5421-5424.

STEFAN, N., M. SARI, T. ECKARDT, T. MICHEL, T. JANE, A. GÖRAN, E. EVA, P. KATARINA, W. MARGARET and G. JAN-ÅKE (2001). "Mechanism of Estrogen Action." Physiological Reviews **81**(4): 1535-1565.

STEIN, B. and M. YANG, X. (1995). "Repression of the Interleukin-6 Promoter by Estrogen Receptor is mediated by NF- κ B and C/EBP β ." Molecular and Cellular Biology **15**(9): 4971-4979.

STENDAHL, M., L. RYDÉN, B. NORDENSKJÖLD, P. JÖNSSON, E., G. LANDBERG and K. JIRSTRÖM (2006). "High progesterone receptor expression correlates to the effect of adjuvant tamoxifen in premenopausal breast cancer patients." Clinical Cancer Research **12**(15): 4614-4618.

SUI, X., K. BRAMLETT, S., M. JORGE, C., D. SWANSON, A., A. VON ESCHENBACH, C. and G. JENSTER (1999). "Specific Androgen Receptor Activation by an Artificial Coactivator." Journal of Biological Chemistry **274**(14): 9449-9454.

SUMAN, S., P. K. SHARMA, G. RAI, S. MISHRA, D. ARORA, P. GUPTA and Y. SHUKLA (2015). "Current perspectives of molecular pathways involved in chronic Inflammation-mediated breast cancer." Biochemical and Biophysical Research Communications.

SUTHERLAND, R., L., R. HALL, E., G. PANG, Y, N., E. MUSGROVE, A. and C. CLARKE, L. (1988). "Effect of medroxyprogesterone acetate on proliferation and cell cycle kinetics of human mammary carcinoma cells." Cancer Research **48**: 5084-5091.

SUTTON, L. M., D. CAO, V. SARODE, K. H. MOLBERG, K. TORGBE, B. HALEY and Y. PENG (2012). "Decreased Androgen Receptor Expression Is Associated With Distant Metastases in Patients With Androgen Receptor-Expressing Triple-Negative Breast Carcinoma." American Journal of Clinical Pathology **138**(4): 511-516.

SZELEI, J., J. JESUS, M. S. ANA, F. L. MARIA and S. CARLOS (1997). "Androgen-Induced Inhibition of Proliferation in Human Breast Cancer MCF7 Cells Transfected with Androgen Receptor." Endocrinology **138**(4): 1406-1412.

TANG, D., S. XU, Q. ZHANG and W. ZHAO (2012). "The expression and clinical significance of the androgen receptor and E-cadherin in triple-negative breast cancer." Medical Oncology **29**(2): 526-533.

TEULINGS, F. A. G., H. A. VAN GILSE, M. S. HENKELMAN, H. PORTENGEN and J. ALEXIEVA-FIGUSCH (1980). "Estrogen, Androgen, Glucocorticoid, and Progesterone Receptors in Progestin-induced Regression of Human Breast Cancer." Cancer Research **40**(7): 2557-2561.

THOMAS, C., G. RAJAPAKSA, F. NIKOLOS, R. HAO, A. KATCHY, C. MCCOLLUM, W., M. BONDESSON, P. QUINLAN, A. THOMPSON, S. KRISHNAMURTHY, F. ESTEVA, J. and J.-Å. GUSTAFSSON (2012). "ER β 1 represses basal-like breast cancer epithelial to mesenchymal transition by destabilizing EGFR." Breast Cancer Research : BCR **14**(6): R148-R148.

THOMAS, M., P. and B. POTTER, V, L. (2013). "The structural biology of oestrogen metabolism." The Journal of Steroid Biochemistry and Molecular Biology **137**(0): 27-49.

TOFT, D., G. SHYAMALA and J. GORSKI (1967). "A receptor molecule for estrogens: studies using a cell-free system." Proceedings of the National Academy of Sciences of the United States of America **57**(6): 1740-1743.

TREMBLAY, A., G. B. TREMBLAY, F. LABRIE and V. GIGUÈRE (1999). "Ligand-Independent Recruitment of SRC-1 to Estrogen Receptor β through Phosphorylation of Activation Function AF-1." Molecular Cell **3**(4): 513-519.

TSANG, J. Y. S., Y.-B. NI, S.-K. CHAN, M.-M. SHAO, B. K. B. LAW, P. H. TAN and M. T. GARY (2014). "Androgen receptor expression shows distinctive significance in ER positive and negative breast cancers." Annals of surgical oncology **21**(7): 2218-2228.

TSUCHIYA, Y., M. NAKAJIMA and T. YOKOI (2005). "Cytochrome P450-mediated metabolism of estrogens and its regulation in human." Cancer Letters **227**(2): 115-124.

UDVARDI, M. K., T. CZECHOWSKI and W. R. SCHEIBLE (2008). "Eleven golden rules of quantitative RT-PCR." Plant Cell **20**(7): 1736-1737.

UEDA, A., K. OKUDA, S. OHNO, A. SHIRAI, T. IGARASHI, K. MATSUNAGA, J. FUKUSHIMA, S. KAWAMOTO, Y. ISHIGATSUBO and T. OKUBO (1994). "NF-kappa B and Sp1 regulate transcription of the human monocyte chemoattractant protein-1 gene." The Journal of Immunology **153**(5): 2052-2063.

VERA-BADILLO, F. E., A. J. TEMPLETON, P. DE GOUVEIA, I. DIAZ-PADILLA, P. L. BEDARD, M. AL-MUBARAK, B. SERUGA, I. F. TANNOCK, A. OCANA and E. AMIR (2013). "Androgen Receptor Expression and Outcomes in Early Breast Cancer: A Systematic Review and Meta-Analysis." Journal of the National Cancer Institute.

VERHOOG, N., J. D., E. JOUBERT and A. LOUW (2007). "Evaluation of the Phytoestrogenic Activity of Cyclopia genistoides (Honeybush) Methanol Extracts and Relevant Polyphenols." Journal of Agricultural and Food Chemistry **55**(11): 4371-4381.

VERKOOIJENA, H., M., C. BOUCHARDYD., V. VINH-HUNGE., E. RAPITID. and M. HARTMAN (2009). "The incidence of breast cancer and changes in the use of hormone replacement therapy: A review of the evidence." Maturitas **64**: 80-85.

VERRAS, M., J. LEE, H. XUE, T.-H. LI, Y. WANG and Z. SUN (2007). "The androgen receptor negatively regulates the expression of c-Met: implications for a novel mechanism of prostate cancer progression." Cancer research **67**(3): 967-975.

VIGNON, F., S. BARDON, D. CHALBOS and H. ROCHEFORT (1983). "Antiestrogenic Effect of R5020, a Synthetic Progestin in Human Breast Cancer Cells in Culture." The Journal of Clinical Endocrinology & Metabolism **56**(6): 1124-1130.

VISSER, K., M. MORTIMER and A. LOUW (2013). "Cyclopia Extracts Act as ER α Antagonists and ER β Agonists, In Vitro and In Vivo." PLoS One **8**(11).

WALTER, P., S. GREEN, G. GREENE, A. KRUST, J. BORNERT, M., J. JELTSCH, M., A. STAUB, E. JENSEN, G. SCRACE and M. WATERFIELD (1985). "Cloning of the human estrogen receptor cDNA." Proceedings of the National Academy of Sciences **82**(23): 7889-7893.

WANG, Q., W. LI, Y. ZHANG, X. YUAN, K. XU, J. YU, Z. CHEN, R. BEROUKHIM, H. WANG, M. LUPIEN, T. WU, M. M. REGAN, C. A. MEYER, J. S. CARROLL, A. K. MANRAI, O. A. JÄNNE, S. P. BALK, R. MEHRA, B. HAN, A. M. CHINNAIYAN, M. A. RUBIN, L. TRUE, M. FIORENTINO, C. FIORE, M. LODA, P. W. KANTOFF, X. S. LIU and M. BROWN (2009). "Androgen Receptor Regulates a Distinct Transcription Program in Androgen-Independent Prostate Cancer." Cell **138**(2): 245-256.

WANG, Y., J. LAM, B., K. LAM, S. L., J. LIU, M. LAM, C., R. HOO, L. C., D. WU, G. COOPER, J. S. and A. XU (2006). "Adiponectin Modulates the Glycogen Synthase Kinase-3 β / β -Catenin Signaling Pathway and Attenuates Mammary Tumorigenesis of MDA-MB-231 Cells in Nude Mice." Cancer Research **66**(23): 11462-11470.

WATSON, C. S., Y.-J. JENG and M. Y. KOCHUKOV (2008). "Nongenomic actions of estradiol compared with estrone and estriol in pituitary tumor cell signaling and proliferation." The FASEB Journal **22**(9): 3328-3336.

WATSONA, C. S., A. M. NORFLEET, T. C. PAPPAS and B. GAMETCHU (1999). "Rapid actions of estrogens in GH 3/B6 pituitary tumor cells via a plasma membrane version of estrogen receptor- α ." Steroids **64**(1): 5-13.

WEATHERMAN, R. V., N. J. CLEGG and T. S. SCANLAN (2001). "Differential SERM activation of the estrogen receptors (ER α and ER β) at AP-1 sites." Chemistry & biology **8**(5): 427-436.

WEATHERMAN, V. R., R. J. FLETTERICK. and T. S. SCANLAN. (1999). "Nuclear-Receptor Ligands and Ligand-Binding Domains." Annual Review of Biochemistry **68**(1): 559-581.

- WELBOREN, W.-J., H. STUNNENBERG, G., F. SWEEP, C, G, J. and P. SPAN, N. (2007). "Identifying estrogen receptor target genes." Molecular Oncology **1**(2): 138-143.
- WELLER, M., A., P. KUPELIAN, A., C. REDDY, A., K. STEPHANS, L. and R. TENDULKAR, D. (2014). "Adjuvant Versus Neoadjuvant Androgen Deprivation With Radiotherapy for Prostate Cancer: Does Sequencing Matter?" Clinical Genitourinary Cancer(0).
- WELSHONS, W., V., M. WOLF, F., C. MURPHY, S. and V. JORDAN, CRAIG. (1988). "Estrogenic activity of phenol red." Molecular and Cellular Endocrinology **57**(3): 169-178.
- WEN-YANG, H., S. GUANG-BIN, H. DAN-PING, N. JASON L and S. GAIL, PRINS. (2012). "Actions of estrogens and endocrine disrupting chemicals on human prostate stem/progenitor cells and prostate cancer risk." Molecular and Cellular Endocrinology **354**(1-2): 63-73.
- WILLIAMS, C., K. EDVARDSSON, S. A. LEWANDOWSKI, A. STROM and J. A. GUSTAFSSON (2007). "A genome-wide study of the repressive effects of estrogen receptor beta on estrogen receptor alpha signaling in breast cancer cells." Oncogene **27**(7): 1019-1032.
- WITZEL, I., M. GRAESER, T. KARN, M. SCHMIDT, R. WIRTZ, D. SCHÜTZE, A. RAUSCH, F. JÄNICKE, K. MILDE-LANGOSCH and V. MÜLLER (2013). "Androgen receptor expression is a predictive marker in chemotherapy-treated patients with endocrine receptor-positive primary breast cancers." Journal of Cancer Research and Clinical Oncology **139**(5): 809-816.
- WONG, C.-W., C. MCNALLY, E. NICKBARG, B. S. KOMM and B. J. CHESKIS (2002). "Estrogen receptor-interacting protein that modulates its nongenomic activity-crosstalk with Src/Erk phosphorylation cascade." Proceedings of the National Academy of Sciences **99**(23): 14783-14788.
- WONG, M., L. and J. MEDRANO, F. (2005). "Real-time PCR for mRNA quantitation." BioTechniques The Journal of Life Science Methods **39**(1): 75-85.
- WOTIZ, H., H., S. CHATTORAJ, C., M. KUDISCH and R. MULLER, ENZIO. (1978). "Impeding Estrogens and the Etiology of Breast Cancer." Cancer Research **38**(11 Part 2): 4012-4020.
- WU, F. and S. SAFE (2007). "Differential activation of wild-type estrogen receptor $\hat{I}\pm$ and C-terminal deletion mutants by estrogens, antiestrogens and xenoestrogens in breast cancer cells." The Journal of Steroid Biochemistry and Molecular Biology **103**(1): 1-9.
- WU, Y., L. SUN, W. ZOU, J. XU, H. LIU, W. WANG, X. YUN and J. GU (2012). "Prosaposin, a regulator of estrogen receptor alpha, promotes breast cancer growth." Cancer Science **103**(10): 1820-1825.

XING, Y., J. P. JIA, X. J. JI and T. TIAN (2013). "Estrogen associated gene polymorphisms and their interactions in the progress of Alzheimer's disease." Progress in Neurobiology **111**(0): 53-74.

XU, J. and Q. LI (2003). "Review of the in Vivo Functions of the p160 Steroid Receptor Coactivator Family." Molecular Endocrinology **17**(9): 1681-1692.

XU, L. L., Y. P. SU, R. LABICHE, T. SEGAWA, N. SHANMUGAM, D. G. MCLEOD, J. W. MOUL and S. SRIVASTAVA (2001). "Quantitative expression profile of androgen-regulated genes in prostate cancer cells and identification of prostate-specific genes." International Journal of Cancer **92**(3): 322-328.

YAGER, J., D. and N. DAVIDSON, E. (2006). "Estrogen Carcinogenesis in Breast Cancer." New England Journal of Medicine **354**(3): 270-282.

YAMEI, Z., Y. CHRISTINA, W. JOE, GRAY., C. KAREN, H. SHANAZ, DAIRKEE. , H. DAN, MOORE., E. URS, E.-C. SERENELLA and C. CHRISTOPHER, BENZ. (2007). "Enhanced NF κ B and AP-1 transcriptional activity associated with antiestrogen resistant breast cancer." BMC Cancer **7**(59).

YANAGAWA, M., K. IKEMOT, S. KAWAUCHI, T. FURUYA, S. YAMAMOTO, M. OKA, A. OGA, Y. NAGASHIMA and K. SASAKI (2012). "Luminal A and luminal B (HER2 negative) subtypes of breast cancer consist of a mixture of tumors with different genotype." BMC research notes **5**(1): 376.

YANG, J., D. SINGLETON, W., E. SHAUGHNESSY, A. and S. KHAN, A. (2008). "The F-domain of estrogen receptor- α inhibits ligand induced receptor dimerization." Molecular and Cellular Endocrinology **295**(1-2): 94-100.

YANG, X., D. L. PHILLIPS, A. T. FERGUSON, W. G. NELSON, J. G. HERMAN and N. E. DAVIDSON (2001). "Synergistic Activation of Functional Estrogen Receptor (ER)- α by DNA Methyltransferase and Histone Deacetylase Inhibition in Human ER- α -negative Breast Cancer Cells." Cancer Research **61**(19): 7025-7029.

YU, H., M. A. LEVESQUE, G. M. CLARK and E. P. DIAMANDIS (1998). "Prognostic value of prostate-specific antigen for women with breast cancer: a large United States cohort study." Clinical cancer research **4**(6): 1489-1497.

YU, Q., Y. NIU, N. LIU, J. Z. ZHANG, T. J. LIU, R. J. ZHANG, S. L. WANG, X. M. DING and X. Q. XIAO (2011). "Expression of androgen receptor in breast cancer and its significance as a prognostic factor." Annals of Oncology **22**(6): 1288-1294.

YUE, W., J. YAGER, D., J.-P. WANG, E. JUPE, R. and R. SANTEN, J. (2013). "Estrogen receptor-dependent and independent mechanisms of breast cancer carcinogenesis." Steroids **78**(2): 161-170.

ZERVOUDIS, S., G. IATRAKIS, E. TOMARA, A. BOTHOU, G. PAPADOPOULOS and G. TSAKIRIS (2014). "Main controversies in breast cancer." World Journal of Clinical Oncology **5**(3): 359-373.

ZHANG, Y., J.-C. CHENG, H.-F. HUANG and P. LEUNG, C, K. (2013). "Homeobox A7 stimulates breast cancer cell proliferation by up-regulating estrogen receptor-alpha." Biochemical and Biophysical Research Communications **440**(4): 652-657.

ZHANG, Z., D. ZHOU, Y. LAI, Y. LIU, X. TAO, Q. WANG, G. ZHAO, H. GU, H. LIAO, Y. ZHU, X. XI and Y. FENG (2012). "Estrogen induces endometrial cancer cell proliferation and invasion by regulating the fat mass and obesity-associated gene via PI3K/AKT and MAPK signaling pathways." Cancer Letters **319**(1): 89-97.

ZHAO, C., K. DAHLMAN-WRIGHT and J.-Å. GUSTAFSSON (2010). "Estrogen Signaling via Estrogen Receptor beta." Journal of Biological Chemistry **285**(51): 39575-39579.

ZHAO, C., D.-W. KARIN and G. JAN-ÅKE (2008). "Estrogen receptor β : an overview and update." Nuclear Receptor Signaling Atlas **6**.

ZILLI, M., A. GRASSADONIA, N. TINARI, A. DI GIACOBBE, S. GILDETTI, J. GIAMPIETRO, C. NATOLI and S. IACOBELLI (2009). "Molecular mechanisms of endocrine resistance and their implication in the therapy of breast cancer." Biochimica et Biophysica Acta (BBA) - Reviews on Cancer **1795**(1): 62-81.

ADDENDUM

Realtime quantitative polymerase chain reaction (qPCR)

Realtime quantitative PCR is a powerful method used for the measurement of gene expression (Celi *et al.* 1994). qPCR is based on the original PCR method that employs forward and reverse primer sequences that are specific for certain target genes. However, qPCR has been modified to involve the use of fluorescent dyes or probes (Pfaffl 2001, Huang *et al.* 2014, Navarro *et al.* 2015). Fluorescent dyes such as SYBR-green are used during qPCR as it binds to the minor groove of the double stranded DNA. The use of SYBR-green allows simultaneous PCR amplification, detection of the PCR product, quantification of the new double stranded DNA being formed and lastly verification of the PCR product using melt curve analysis (Morrison and Weiss 1998, Phillips *et al.* 2009). However, there are certain factors that should be considered when conducting qPCR such as making sure that the reference gene does not give any expression under the experimental conditions as well as using RNA that is of good quality (Bustin 2002, Udvardi *et al.* 2008).

To assess gene expression, high quality, intact RNA has to be isolated to ensure that experimental results obtained are reliable as poor quality RNA can compromise results. RNA quality was assessed by determination of the optical density (OD) at 260 nm (nucleic acids) and the 280 nm (proteins), followed by analysis on a 1% denaturing formaldehyde agarose gel. The latter is to analyse the quality and integrity of RNA isolated from the cells as the 260 nm OD reading can be compromised by the presence of genomic DNA (Sambrook *et al.* 1989). An OD 260/280 ratio between 1.9 and 2.1 indicates that the RNA is pure (Sambrook *et al.* 1989). As shown in figure A1, the intact total RNA isolated from the MDA-MB-231 cells showed sharp, clear 28S and 18S RNA bands. As the 28S RNA band is approximately twice the intensity of the 18S RNA band, it is a good indication that the RNA is intact (Krebs *et al.* 2009). If the 28S and 18S RNA bands appearance as a smear, it is an indication that the RNA is degraded, while genomic DNA does not migrate through the gel because of its size (Sambrook *et al.* 1989).




Figure A1. A representative 1% denaturing formaldehyde agarose gel showing the intact total RNA isolated from MDA-MB-231 cells. Total RNA was isolated from MDA-MB-231 cells transiently transfected with 2.5 ng pGL2basic, ER α or ER β , as described in section 2.6 of Chapter 2. A total of 1 μ g RNA was electrophoresed on a 1% agarose gel and visualised by ethidium bromide staining.

The intact RNA was then used to synthesize complementary DNA (cDNA) as described in section 2.6 of Chapter 2.

Determination of primer pair efficiency

To calculate the relative expression of the genes of interest by the method of (Pfaffl 2001), the primer efficiencies have to be determined. Primer efficiency is generally assumed to be two, since the product is doubled after every PCR cycle (Pfaffl 2001). To determine the primer efficiency, a serial dilution of a single cDNA sample was made, and each dilution was analysed in triplicate. A no template control was added, to determine if any contamination or primer self-amplification occurred. The samples were then analysed using the qPCR protocol described in section 2.7 of Chapter 2. A standard curve was generated using the StepOne™ software by plotting the cycle number (CT) (Y-axis) for each dilution against the log of the cDNA concentration (X-axis) (figure A2). The slope obtained from the standard curve is then used to

calculate the exponential amplification value (E), which is the primer efficiency for each primer pair, by using the following equation (Pfaffl 2001).

$$E = 10^{(-1/\text{slope})} \quad (\text{Equation 1})$$

The primer efficiencies were 1.96, 1.94 and 2.12 for PR, PSA and GAPDH, respectively.




Figure A2 continues on next page.




Figure A2. Representative standard curve showing cycle number (CT) versus the log DNA concentration for the (A) PR, (B) PSA and (C) GAPDH genes, respectively. The slope obtained from the each graph was used to calculate the primer efficiency.

Melting curve analysis

Melting curves are used as a way of optimization to confirm primer specificity, to determine whether there are any non-specific products and primer dimers. Primer dimers are indicated by the presence of small peaks that protrude from the product peak, while non-specific products can be indicated by the presence of peaks that deviate from the melting temperature (T_m) line (indicated by the vertical blue line in figure A3). During the PCR reaction SYBR green forms complexes with the double stranded DNA products. Therefore the fluorescence measured using SYBR green is the measurement of the sum of the relative levels of all products formed during the reaction. It is very important to ensure that the measured fluorescence is of a single product. This is done by increasing the temperature of the samples by 0.2°C per second until the product's denaturation temperature of 95°C is passed. The melting curve of a PCR product is dependent on the GC/AT content, length and sequence of the PCR product (Ririe *et al.* 1997). A melting curve is generally a plot of fluorescence as a function of temperature, and one product is indicated by one peak of high amplitude on the melting curve (Ririe *et al.* 1997). Examples

of melting curves for PR, PSA and GAPDH, respectively are shown below in figure A3, and a 2% agarose gel was used to further confirm that a single amplicon of the correct size was amplified (figure A4 (A-C)).




Figure A3 continues on next page.




Figure A3 legend on next page.

Figure A3. Representative melting curves for the PR, PSA and GAPDH genes, respectively. MDA-MB-231 cells were transfected with (A and C) pSG5-ER β and subsequently treated with 1 nM E₂, while MDA-MB-231 cells transfected with (B) pSVAR_o were treated with 1 nM DHT. The red line represents samples treated with 1 nM E₂, the blue line indicates samples treated with 1 nM DHT and the grey line is the no template control.




Figure A4 legend on next page.

Figure A4 Representative agarose gels showing qPCR products of (A) PR, (B) PSA and (C) GAPDH. Human MDA-MB-231 breast cancer cells were transfected with 2.5 ng (A) and (C) pSG5-hER β or (B) pSVAR_O using X-tremeGENE HP DNA transfection reagent according to the manufacturer's instructions for 24 hours. Cells were subsequently treated with 1 nM E₂ (PR and GAPDH) or DHT (PSA) for 12 hours. Total RNA was isolated, cDNA synthesized and mRNA expression levels of PR, PSA and GAPDH were determined by qPCR. The products were electrophoresed on a 2% agarose gel and visualized using ethidium bromide staining. Lane 1 represents the no template control, while lane 2 represents the single PCR amplicon of the correct size.

Determination of relative gene expression values

Having determined that a single amplicon of the correct size had been amplified, the relative expression (R) levels of the gene of interest were calculated using the mathematical model below (Pfaffl 2001), and expressed relative to the expression levels of the vehicle control (EtOH)

$$R = \frac{(E_{\text{target gene}})^{\Delta CP(\text{control-sample})}}{(E_{\text{reference gene}})^{\Delta CP(\text{control-sample})}} \quad (\text{Equation 2})$$

The crossing point (CP) is the point in the PCR cycle where the target amplicon is first detected, and can also be referred to as the cycle number (C_t). C_t is used in ABI Prism[®] literature (Applied Biosystems, USA), while CP is used in LightCycler[®] literature (Roche Applied Science, USA) (Wong and Medrano 2005). The ΔCP is calculated for both the target and reference genes (GAPDH), by subtracting the CP value of the treated sample from the CP value of the vehicle control (EtOH) (Pfaffl 2001). A ratio of one indicates that there is no difference between the treated sample and that of the vehicle control. Results of experiments are graphically presented by plotting the cycle number versus the log concentration of the amplified cDNA.

Determining the optimal time for gene expression

MDA-MB-231 cells were transfected with 2.5 ng of either ER α , ER β , AR or the pGL2 basic empty vector for 24 hours. The cells transfected with ER α , ER β or pGL2 basic were treated with 0.1% EtOH or 1 nM E₂, while cells transfected with the AR were treated with 0.1% EtOH or 1 nM DHT for 2, 6, 12 or 24 hours. As shown in figure A5 A, E₂ did not upregulate the expression of the PR gene at any of the four time points in cells transfected with the empty vector. In contrast, cells transfected with ER α or ER β showed significant E₂-induced upregulation of the PR gene (figure A5 B and C). Similarly, significant DHT-induced upregulation of the PSA gene was seen in cells transfected with the AR (figure A5 D). As good induction of both the PR (4.5-fold in the presence of ER α ; 3.5-fold in the presence ER β) and PSA genes (2.1-fold in the presence of the AR) were observed at 12 hours, this treatment time was chosen for all subsequent experiments.




Figure A5 legend on next page.

Figure A5. Optimal gene expression for human PR and PSA. Human MDA-MB-231 breast cancer cells were transfected with 2.5 ng (A) pGI2 basic or (B) pSG5-hER α or (C) pSG5-hER β or (D) pSVAR_o using X-tremeGENE HP DNA transfection reagent according to the manufacturer's instructions for 24 hours. Cells were subsequently treated with 0.1% (v/v) EtOH (control) or 1 nM E₂ (PR) or DHT (PSA) for 2, 6, 12 and 24 hours. Total RNA was isolated, cDNA synthesized and mRNA expression levels of PR, PSA and GAPDH were determined by QPCR. GAPDH was used as the internal standard and the ratio of PR or PSA mRNA expression/GAPDH mRNA of all samples were calculated relative to the sample treated with 0.1% (v/v) EtOH. Results indicate the PR mRNA expression at 1 nM E₂ via (A) pGI2 basic or (B) ER α or (C) ER β . (D) Results indicate the PSA mRNA expression at 1 nM DHT via AR.

